



CAPTAN 65

SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE CONTROL — DYE ADDED

ACTIVE INGREDIENT:	65
Captan*	
INERT INGREDIENTS	35
*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide	100%

CAUTION

Keep Out of Reach of Children

CAUSES SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED OR INHALED. Avoid contact with skin, eyes, or clothing. Avoid breathing dust — wear respirator. Wash thoroughly after handling. Do not store or use near feed or food. In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Wash skin with soap and water.

PROTECT FROM EXCESSIVE HEAT — STORE IN A COOL DRY PLACE

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES.

CAPTAN 65 SEED PROTECTANT is a microfine wettable powder especially formulated for the treatment of seed, prior to storage and planting, to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

COMPATIBILITY: Captan 65 can be used in combination with most commonly used seed protectants, both insecticides and fungicides as well as fertilizers. Do not use with strongly alkaline materials.

PLANT SAFETY: At the recommended dosage Captan 65 is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated, giving a superior result as compared to results from other seed fungicides.

GENERAL USE PRECAUTIONS

Read all directions before using and use only as specified on this label.
Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.
Dispose of excess treated seeds by burial.
IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled.

TREATED SEED — DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

DIRECTIONS FOR USE

SLURRY TREATMENT

RECOMMENDATIONS

AMOUNT OF CAPTAN 65 TO USE ON SEEDS

SEED	SLURRY	SEED	SLURRY
ALFALFA, CLOVER, ESPERDEZA, TREFOIL	6 1/2 oz./100 lb.	CRUCIFERS: Mustard, Radishes, Rape, Turnips	1 1/4 oz./100 lb.
BARLEY, OATS, WHEAT	1 1/2 oz./bu.	FLAX	1 1/2 oz./bu.
BEANS (Dry), Snap, Lima, Cowpeas, Peas, Soybeans	1 1/4 oz./bu.	GRASSES	6 1/2 oz./100 lb.
BEETS (Garden)	9 1/2 oz./100 lb.	MUSKMELON, PUMPKIN, SQUASH, WATERMELON	1 1/2 oz./100 lb.
BLUE GRASS	6 1/2 oz./100 lb.	ONIONS (Pelleting)	1 1/4 lb./1 lb.
CANTALOUPE, CUCUMBER	2 1/2 oz./100 lb.	PEPPER (California Wonder)	2 1/2 oz./100 lb.
COLE CROPS (Broccoli, Brussels, Sprouts, Cabbage, Cauliflower)	1 1/4 oz./100 lb.	SORGHUM (Hull-less)	3 1/2 oz./100 lb.
CORN — FIELD	1 1/2 oz./bu.	SPINACH	4 1/2 oz./100 lb.
SWEET CORN	3 oz./100 lb.	SUGAR BEETS — West	4 1/2 oz./100 lb.
COTTON — ACID DELINTED, REGINNED, FUZZY	2 1/2 oz./100 lb. 3 1/2 oz./100 lb. 3 1/2 oz./100 lb.	East	9 1/2 oz./100 lb.
		SWISS CHARD	9 1/2 oz./100 lb.

GLADIOLUS BULBS: Prevention of bulb rot — Use 1 1/2 oz. in 1 gal. of water (1 lb. Captan 65 to 9 gals. of water). Soak bulbs in mixture 20 to 30 minutes.

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose. This material, except those which are contained on Stauffer's label.

EPA Reg. No. 476-1970

LBS. NET

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

