

CAPTAN-STREPTOMYCIN 5-01

POTATO SEED PIECE PROTECTANT

AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Captan: N-[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide	5.00%
Streptomycin*	0.01%

INERT INGREDIENTS	94.99%
	100.0%

Not licensed for use in countries where valid patents still exist.


*Streptomycin from streptomycin sulfate (agricultural grade)
Contains 100 ppm. streptomycin.

CAUTION

Keep Out of Reach of Children

Harmful If Swallowed

- May cause skin irritation.
- Avoid contact with skin and eyes
- Avoid breathing dust.
- Wash with soap and water after exposure
- Avoid contamination of food and feed



RECOMMENDATIONS

This material is especially formulated for potato seed piece treatment.

POTATOES (seed pieces): seed rot and damp-off, verticillium wilt, brown eye disease. Dust at 15 ounces per bushel cut seed pieces (24 ounces or 1½ lb. to 100 lb. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored 2 or 3 days in open crates before bagging.

Further specific information on this subject may be obtained from state agricultural experiment stations or the State Agricultural Extension Service.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability, and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

- PROTECT FROM EXCESSIVE HEAT.
- STORE IN A COOL DRY PLACE.
- BURN EMPTY BAG.
- KEEP OUT OF SMOKE.

TOXIC TO FISH. KEEP OUT OF ANY BODY OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES

U.S.D.A. Reg. No. 476-197.

ARC 710211

CAPTAN-STREPTOMYCIN

5-01

POTATO SEED PIECE PROTECTANT AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Captan: N-[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide	5.00%
Streptomycin*	0.01%

INERT INGREDIENTS

	94.99%
	100.0%

Not licensed for use in countries where valid patents still exist.

*Streptomycin from streptomycin sulfate (agricultural grade)
Contains 100 ppm. streptomycin.

RECOMMENDATIONS

This material is especially formulated for potato seed piece treatment.

POTATOES (seed pieces): seed rot and damp-off, verticillium wilt, brown eye disease. Dust at 15 ounces per bushel cut seed pieces (24 ounces or 1½ lb. to 100 lb. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored 2 or 3 days in open crates before bagging.

Further specific information on this subject may be obtained from state agricultural experiment stations or the State Agricultural Extension Service.

CAUTION

Keep Out of Reach of Children

Harmful If Swallowed


May cause skin irritation

Avoid contact with skin and eyes.

Avoid breathing dust.

Wash with soap and water after exposure.

Avoid contamination of food and feed.



NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability, and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

PROTECT FROM EXCESSIVE HEAT

STORE IN A COOL DRY PLACE.

BURN EMPTY BAG.

KEEP OUT OF SMOKE.

TOXIC TO FISH. KEEP OUT OF ANY BODY OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTE.