

CAPTAN-SULFUR 5-50 DUST

AN AGRICULTURAL FUNGICIDE

Active Ingredients:

Captan: N-[(trichloromethyl)thio]4-cyclohexene-1,2-dicarboximide . . . 5%

Sulfur 50%

Inert Ingredients 45%

U.S. Pat. Nos. 2,553,770-2,553,771

Not licensed for use or export outside of the United States, its territories and possessions, Canada or Mexico.

CAUTION

Keep Out of Reach of Children

AVOID INHALING DUST.

AVOID CONTACT WITH SKIN.

AVOID CONTAMINATION OF FEED AND FOOD.

~~Keep Out of Reach of Children~~

OBSERVE THE FOLLOWING USE PRECAUTIONS

USE ONLY ON CROPS RECOMMENDED

Do not use on severe fruit and leaf injury to sulfur sensitive crops. Do not use on apples, pears, Anjou and Comice pears, cranberries, cucurbits (cucumbers, melons, squash), walnuts or to other sensitive plants. Do not use on any crop unless local use has proved safe. Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures are likely to exceed 90 F.

Insecticidal and acaricidal residues on feed and forage crops will not exceed tolerance established by the Federal Food and Drug Administration, use only at intervals which do not interfere with harvest. Do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or livestock or other animals, feed, fiber or pasture crops. The grower is responsible for residues on crops as well as for damages caused by drift to property to that of others. Consult State Agricultural Experiment Station or the State Agricultural Extension Service, as the timing, number, and intervals of applications needed will vary with local conditions.

This product is not to be used on grapes, citrus, and pears.

DIRECTIONS FOR USE

Unless otherwise specified, apply at first sign of infestation and infection by ground equipment or aircraft. Unless otherwise stated repeat at 5-10 day intervals as needed to maintain control.

Maximum permissible rates per acre of the formulation are given in parentheses after each crop.


RECOMMENDATIONS and DIRECTIONS FOR USE

STRAWBERRIES - Botrytis, powdery mildew. - Apply ~~30~~ ²⁰ lb. per acre when new growth starts in the spring before fruit forms. Repeat dustings at 7-10 day intervals. (~~30~~ ²⁰ lb.)

NOTICE: Stauffer Chemical Company makes no warranty of merchantability or any other express or implied warranty concerning this material. It shall not be held responsible for personal injury, property damage, or other losses resulting from the handling, storage or use of this material. The buyer assumes all risk and liability resulting from such handling, storage or use.

Made in U.S.A.
Stauffer Chemical Company
New York

67-1000-101
1974



50 LB. NET

BURN EMPTY BAG - KEEP OUT OF SMOKE