

CAPTAN-SULFUR 7.5-50 DUST

AN AGRICULTURAL FUNGICIDE

Active Ingredients:

Captan: N-[(trichloromethyl) thio] 4-cyclohexene-1,2-dicarboximide... 7.5

Sulfur... 50.0

Inert Ingredients

42.5

U.S. Pat. Nos. 2,553,770 - 2,553,771

Not licensed for use or export outside of the United States, its territories and possessions, Canada or Mexico.

CAUTION

Keep Out of Reach of Children

AVOID INHALING DUST

AVOID CONTACT WITH SKIN.

AVOID CONTAMINATION OF FEED AND FOOD.

~~Keep Out of Reach of Children~~

OBSERVE THE FOLLOWING USE PRECAUTIONS

USE ONLY ON CROPS RECOMMENDED

Do not use on severe fruit and leaf injury to sulfur-sensitive crops. Do not apply to peaches, Anjou and Comice pears, strawberries, cucurbits (cucumbers, melons, squash, pumpkins), walnuts or to other sensitive plants. Do not use on any crop unless local use has proved safe. Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures are likely to exceed 90° F.

Do not use if post-harvest residues on food and storage crops will not meet the tolerance established by the Federal Food and Drug Administration. Do not use on crops in States and territories, and do not apply close to harvest time, or to fruit to be applied or allowed to left to areas occupied by uninfected plants. Do not use on animals or on pasturing food, fiber or pasture crops. The grower is responsible for use of this material as well as for damages caused by drift from application to third parties. Consult State Agricultural Experiment Stations or the State Agricultural Extension Service, as the timing, number, and amount of applications needed will vary with local conditions.

This product is toxic to fish. Keep out of lakes, streams and ponds.

DIRECTIONS FOR USE


Unless otherwise specified, apply at first sign of infestation and infection by ground equipment or aircraft. Unless otherwise stated repeat at 5-10 day intervals as needed to maintain control.

Maximum permissible rates per acre of the formulation are given in parentheses after each crop.

RECOMMENDATIONS and DIRECTIONS FOR USE

STRAWBERRIES - Botrytis, powdery mildew. Apply 30 lb. per acre when new growth starts in the spring before fruiting. Repeat at 10-14 day intervals. (38 lb.)

NOTICE: Stauffer Chemical Company makes no warranty of merchantability or any other express or implied warranty concerning this material. It will not be held responsible for personal injury, property damage, or other loss resulting from the handling, storage, or use of this material. The buyer assumes all risk and liability resulting from use of this material.



50 LB. NET

BURN EMPTY BAG KEEP OUT OF SMOKE