

CAPTAN-THIODAN* 5-2 DUST

AGRICULTURAL INSECTICIDE AND FUNGICIDE

ACTIVE INGREDIENTS

CAPTAN: N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide 5.0%

Endosulfan* 2.0%

INERT INGREDIENTS: 93.0%

100.0%

Not licensed for use in countries where valid patents exist.

*Hexachloro-hexahydro-methano-2,4,3-benzodioxathiepin-oxide

Thiodan is a registered trademark of Canadian Hoechst Ltd. U.S. Patent No. 2,799,685

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN

Avoid breathing dust

Avoid contact with skin, eyes or clothing

In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention

Wash with soap and water after handling and before eating or smoking; wear clean clothing.

Avoid contamination of feed and food.

OBSERVE FOLLOWING USE PRECAUTIONS

Observe use instructions given on the label for specific crops.

Be aware that pesticide residues on food and forage crops will be present. This product is registered by the Federal Food and Drug Administration; use only at the maximum tolerance level specified, and do not apply or set to harvest than specified. Do not apply to water or drift to water bodies by any method, human or beneficial animals, or to land, or to land, or to water bodies. The grower is responsible for residues on crops or on water bodies caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as to timing, number and rate of application, and for use with special conditions.

Toxic to fish and wildlife. Do not apply where any body of water may be contaminated by direct application or runoff. Birds feeding on treated areas may be killed.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

BURN EMPTY BAG—KEEP OUT OF SMOKE

DIRECTIONS FOR USE

Apply when a disease is causing ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops.

For the control of disease persisting after the prescribed time limits have expired, an effective fungicide should be used alone, in a manner which will not leave residue in excess of established tolerance.

CAUTION: Workers entering treated area, the day of application, should wear protective clothing.

RECOMMENDATIONS

CUCUMBER, PUMPKIN, SUMMER & WINTER SQUASH: Angular leaf spot, anthracnose, downy mildew, aphids, cucumber beetle, striped cabbage flea beetle, squash vine borer. Apply 40-50 lb. per acre. For squash vine borer control, apply weekly to flower buds, stems and vines beginning when moths first appear. May be applied up to day of harvest.

POTATOES: Early & late blight, aphids, European corn borer, Colorado potato beetle, false chinch bug, flea beetle, green stink bug, leafhopper, leaf-footed plant bug, potato psyllid, potato tuberworm, three-lined potato beetle, Southern armyworm. Apply 40-55 lb. per acre. Repeat as needed.

STRAWBERRIES: Botrytis rot, seed-wasp, spotted wilt, strawberry aphid. Apply 35-50 lb. per acre. Do not apply within 4 days of picking. Do not apply at intervals of less than 10 days or more than twice during any 30-day period if fruit is present.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

50 LB. NET

476-1981
 ACCEPTED
 JUN 19 1970
 THE FEDERAL BUREAU OF INVESTIGATION
 U.S. DEPARTMENT OF JUSTICE
 WASHINGTON, D.C. 20535

USDA REG. NO. 476-1981