

Feb 10 1991 496-1944



Dyfonate[®]

5-G

50 POUNDS NET

ACTIVE INGREDIENT:
Chloro 1,5-bis(phenyl-ethyl)-phosphorodithioate 50.0
INERT INGREDIENTS: 95.0
100.0

GRANULAR SOIL INSECTICIDE
(Contains 5 Pounds Active Ingredient Per 100 Pounds)

U. S. Patent No. 2,988,474)
READ ALL DIRECTIONS BEFORE USING
WARNING!

KEEP OUT OF REACH OF CHILDREN
HARMFUL OR FATAL IF SWALLOWED

SKIN CONTACT OR INHALATION OF DUSTS
MAY BE HARMFUL

DO NOT GET ON SKIN, IN EYES, ON CLOTHING,
AFTER BREATHING DUST. WEAR GLOVES WHEN
HANDLING. IN CASE OF CONTACT, WASH IM-
MEDIATELY WITH SOAP AND WATER. ALWAYS
WASH THOROUGHLY AFTER HANDLING.

NOTE TO PHYSICIANS: Exposure may cause
allergic reaction. Antidote: Atropine. Antidotal
to PAM is also anticholinergic. Antidote: Atropine.

- OBSERVE THE FOLLOWING USE PRECAUTIONS**
- Do not apply to any material that has been treated with a similar product.
 - Do not apply to any material that has been treated with a similar product.
 - Do not apply to any material that has been treated with a similar product.
 - Do not apply to any material that has been treated with a similar product.

- SOIL INSECTS CONTROLLED**
- Cabbage Maggot** (*Delia floralis*)
Garden Symphylan (*Symphyla*)
Northern Corn Rootworm (*Diabrotica*)
Onion Maggot (*Delia floralis*)
Southern Corn Rootworm (*Diabrotica*)
Western Corn Rootworm (*Diabrotica*)
Wireworms:
 Corn Wireworm (*Cryptus*)
 Pacific Coast Wireworm (*Cryptus*)
 Southern Potato Wireworm (*Cryptus*)
 Sugar Beet Wireworm (*Cryptus*)
 Tobacco Wireworm (*Cryptus*)
Western Field Wireworm (*Cryptus*)
Wheat Wireworm (*Cryptus*)

DIRECTIONS FOR USE AND RECOMMENDATIONS

Asparagus: (California Only) For control of Garden Symphylans, apply 60 to 200 pounds per acre, broadcast. Incorporate into the soil with standard cultivation equipment. Do not harvest for 18 months.

Cole Crops: (For Seed Only) For control of Cabbage Maggot prior to and at the time of seeding or transplanting. Prior to seeding or transplanting, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Radish: For control of Cabbage Maggot prior to and at the time of seeding or transplanting. Prior to seeding or transplanting, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Field, Sweet and Pop Corn: For control of Wireworms, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Irish Potatoes: For control of Wireworms, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Bulb Onions: For control of Wireworms, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Directions and Recommendations (continued)

Peanuts: For control of Rootworms, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Sugar Beets: For control of Wireworms, apply 80 pounds per acre, broadcast, and incorporate into the soil by disking.

Sweet Potatoes: For control of Wireworms, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

Tobacco: For control of Wireworms, apply 40 pounds per acre, broadcast, and incorporate into the soil by disking. Do not use treated crops for food or feed purposes.

BEFORE USING — STUDY SAFETY DIRECTIONS

Do not use this material in or near water. Do not use this material in or near water. Do not use this material in or near water.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

STAUFFER CHEMICAL COMPANY
New York, N.Y. 10017

STAUFFER CHEMICAL COMPANY

NEW YORK, N.Y. 10017