

476-1995

12-14-70

leaf 3



Dyfonate®

ACTIVE INGREDIENT:

O-ethyl S-phenyl ethylphosphonodithioate 10.0%

INERT INGREDIENTS: 90.0%

100.0%

GRANULAR SOIL
 (Contains 10 Pounds Active)

(U. S. Patent No. 2,988,474)

READ ALL DIRECTIONS BEFORE USING

WARNING!

KEEP OUT OF REACH OF CHILDREN

HARMFUL OR FATAL IF SWALLOWED

 SKIN CONTACT OR INHALATION OF DUSTS
 MAY BE HARMFUL

DO NOT GET ON SKIN, IN EYES, ON CLOTHING, AVOID BREATHING DUST, WEAR GLOVES WHEN HANDLING. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. ALWAYS WASH THOROUGHLY AFTER HANDLING.

NOTE TO PHYSICIANS: Exposure may cause cholinesterase inhibition. Atropine is Antidotal. 2-PAM also is antidotal when administered early and in conjunction with atropine.

SOIL INSECTS CONTROLLED
Cabbage Maggot (*Hylemya Brassicae*)

Garden Symphylan (*Scutigereilla Immaculata*)

Northern Corn Rootworm (*Diabrotica Longicornis*)

Onion Maggot (*Hylemya Antiqua*)

Southern Corn Rootworm

 (*Diabrotica Undecimpunctata Howardi*)

Western Corn Rootworm (*Diabrotica Virgifera*)

Wireworms:
Corn Wireworm (*Melanotus Communis*)

Pacific Coast Wireworm (*Limonius Canus*)

Southern Potato Wireworm (*Conoderus Falli*)

Sugar Beet Wireworm (*Limonius Californicus*)

Tobacco Wireworm (*Conoderus Vespertinus*)

Western Field Wireworm (*Limonius Infuscatus*)

Wheat Wireworm (*Agriotes Mancus*)

Ctenicera Lobata
Dalopius Pallidus
OBSERVE THE FOLLOWING USE PRECAUTIONS

- Do not apply in any manner not specified on this label.
- Not for use or storage in or around the home.
- Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops.
- This product will kill fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed.

DIRECTIONS FOR USE AND RECOMMENDATIONS

Asparagus: (California Only) For control of Garden Symphylans, apply 30 to 100 pounds per acre, broadcast. Incorporate into the soil with standard cultivation equipment. Do not harvest for 18 months.

Cole Crops: (For Seed Only) For control of Cabbage Maggot prior to and at the time of seeding or transplanting. Prior to seeding or transplanting, apply 20 pounds per acre, broadcast, and incorporate into the soil by discing. Do not use treated crops for food or feed purposes.

Field, Sweet and Pop Corn: For control of Northern and Western Corn Rootworms, apply 7½ to 10 pounds per acre in a 6 to 8-inch band at time of planting over corn rows 38-40 inches apart. Incorporate into the top ½ to 1 inch of soil by making application ahead of press wheels, or by use of special covering devices, or by dragging a short length of chain behind the press wheels.

For control of Wireworms in corn, apply 40 pounds per acre, broadcast, prior to planting and incorporate into the soil by discing.

For control of Symphylans in corn, apply 20 pounds per acre, broadcast, prior to planting and incorporate into the soil by discing.

Irish Potatoes: For control of Wireworms, apply 40 pounds per acre. Apply broadcast and incorporate into the soil by discing prior to planting.

Bulb Onions: For control of Onion Maggot, apply as a furrow treatment at time of seeding. Based on a 20' row spacing apply at a rate of 10 lbs. per acre (10 lbs. per 26,109 ft. of row, 1 lb. per 2,610 ft. of row (¾ lb. per 1,000 ft. of row). May be used on all soil types.

Peanuts: For control of Rootworms, apply 20 pounds per acre in an 18-inch band over the peanut rows 36 inches apart. Apply just prior to pegging and incorporate lightly into the soil with cultivation equipment.

(Continued Next Column)

STAUFFER CHEMICAL COMPANY

2 of 3

50 POUNDS NET

10-G

INSECTICIDE
Ingredient Per 100 Pounds)

Directions and Recommendations (continued)

Radish: For control of Cabbage Maggot prior to and at time of seeding or transplanting, prior to seeding or transplanting, apply 20 pounds per acre, broadcast, and incorporate into the soil by discing.

Sugar Beets: For control of Wireworms, apply 40 pounds per acre broadcast prior to planting and incorporate into the soil by discing.

Sweet Potatoes: For control of wireworms in the southeastern states, apply 20 pounds per acre broadcast and soil incorporate by discing prior to planting. For control of Wireworms in other geographical locations, apply 40 pounds per acre broadcast.

Tobacco: For control of Wireworms as listed above, apply 10 to 20 pounds per acre, broadcast, prior to planting, and incorporate into the soil by discing.

BEFORE USING — STUDY SAFETY DIRECTIONS

Completely empty the contents of bag and bury unused chemical at least 18 inches deep in an isolated location away from water supplies. Burn bag immediately and stay out of smoke.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-1995

ARC701012
OOM701019MW



NEW YORK, NY 10017

