

PLEASE NOTE

This image contains more than one label approved for this product on this date.

Stauffer
CHEMICALS

Dyfonate 10-G

50

ACCEPTED
476-1995
JUN 2 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRA-
TION UNDER NO. SUBJECT
TO ATTACHED COMMENTS

ACTIVE INGREDIENT:

O-ethyl S-phenyl ethylphosphonodithioate 10.0%
INERT INGREDIENTS: 90.0%
100.0%

(U. S. Patent No. 2,988,474)

READ ALL DIRECTIONS BEFORE USING

WARNING!

KEEP OUT OF REACH OF CHILDREN

HARMFUL OR FATAL IF SWALLOWED

SKIN CONTACT OR INHALATION OF DUSTS
MAY BE HARMFUL

DO NOT GET ON SKIN, IN EYES, ON CLOTHING.
AVOID BREATHING DUST. WEAR GLOVES WHEN
HANDLING. IN CASE OF CONTACT, WASH IM-
MEDIATELY WITH SOAP AND WATER. ALWAYS
WASH THOROUGHLY AFTER HANDLING.

NOTE TO PHYSICIANS: Exposure may cause
cholinesterase inhibition. Atropine is Antidotal.
2-PAM also is antidotal when administered early
and in conjunction with atropine.

SOIL INSECTS CONTROLLED

Cabbage maggot (*Hylemya brassicae*)
Garden symphylan (*Scutigera immaculata*)
Northern corn rootworm (*Diabrotica longicornis*)
Onion maggot (*Hylemya antiqua*)
Southern corn rootworm
(*Diabrotica undecimpunctata howardi*)
Western corn rootworm (*Diabrotica virgifera*)
Wireworms:
Corn wireworm (*Melanotus communis*)
Pacific coast wireworm (*Limonius canus*)
Southern potato wireworm (*Conoderus falli*)
Sugar beet wireworm (*Limonius californicus*)
Tobacco wireworm (*Conoderus vespertinus*)
Western field wireworm (*Limonius infuscatus*)
Wheat wireworm (*Agriotes mancus*)
Ctenicera lobato
Dalopius pallidus

OBSERVE THE FOLLOWING USE PRECAUTIONS

- Do not apply in any manner not specified on this label.
- Not for use or storage in or around the home.
- Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops.
- This product will kill fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed.

GRANULAR SOIL INSECTICIDE
(Contains 10 Pounds Active Ingredient Per 100 Pounds)

DIRECTIONS FOR USE AND RECOMMENDATIONS

Asparagus: (California Only) For control of Garden Symphylans, apply 30 to 100 pounds per acre, broadcast. Incorporate into the soil with standard cultivation equipment. Do not harvest for 18 months.

Cole Crops: (For Seed Only) For control of Cabbage Maggot and Garden Symphylans prior to and at the time of seeding or transplanting. Prior to seeding or transplanting, apply 20 pounds per acre, broadcast, and incorporate into the soil by discing. Do not use treated crops for food or feed purposes.

Corn (Field, Sweet and Pop Corn):

A. Band Application for Control of Northern and Western Corn Rootworms:

Apply 7-1/2 to 10 lbs. Dyfonate 10:G per acre in a 6 to 8 inch band over corn rows 38 to 40 inches apart. This is equivalent to 9 to 12 oz. of Dyfonate 10:G per 1,000 linear feet of row. Use this same rate per linear foot of row for any row spacing.

1. **Application at Planting:** Incorporate into the top 1/2 to 1 inch of soil by making application ahead of the press wheels, or by dragging a short length of chain behind the press wheels.

OR
2. **Application at Cultivation:** Cover the treated band with 2-3 inches of soil by making application immediately ahead of disc hillers or cultivation equipment. Make application in May or June when activity of rootworm larvae is first observed. Do not apply within 45 days of harvest.

B. Broadcast Application for Control of Symphylans and Wireworms:

Apply 40 pounds (for Wireworms) or 20 pounds (for Symphylans) Dyfonate 10:G per acre, prior to planting and incorporate into the soil by discing.

Bulb Onions: For control of Onion Maggot, apply as a furrow treatment at time of seeding. Based on a 20" row spacing apply at a rate of 10 lbs. per acre (10 lbs. per 26,109 ft. of row, 1 lb. per 2,610 ft. of row; 3/8 lb. per 1,000 ft. of row). May be used on all soil types.

(Continued Next Column)

Directions and Recommendations (continued)

Peanuts: For control of Rootworms, apply 20 pounds Dyfonate 10:G per acre in an 18-inch band over the peanut rows 36 inches apart. Apply just prior to pegging and incorporate lightly into the soil with cultivation equipment.

Potatoes (Irish): For control of Garden Symphylans and Wireworms, apply 40 pounds (for Wireworms) or 20 pounds (for Symphylans) Dyfonate 10:G per acre, broadcast, prior to planting and incorporate into the soil by discing.

Radish: For control of Cabbage Maggot and Garden Symphylans prior to and at time of seeding or transplanting. Prior to seeding or transplanting, apply 20 pounds per acre, broadcast, and incorporate into the soil by discing.

Sugar Beets: For control of Wireworms and Garden Symphylans, apply 40 lbs. per acre broadcast (for wireworms) or 20 lbs. per acre broadcast (for symphylans) prior to planting and incorporate into the soil by discing.

Sweet Potatoes: For control of wireworms in the southeastern states, apply 20 pounds per acre broadcast and soil incorporate by discing prior to planting. For control of Wireworms in other geographical locations, apply 40 pounds per acre broadcast.

Table Beets: For control of Cabbage Maggot and Garden Symphylans, apply 20 pounds per acre, broadcast, prior to planting. Incorporate into the soil by discing.

Tobacco: For control of Wireworms as listed above, apply 10 to 20 lbs. per acre, broadcast, prior to bedding and transplanting, and incorporate into the soil by discing.

BEFORE USING — STUDY SAFETY DIRECTIONS

Completely empty the contents of bag and bury unused chemical at least 18 inches deep in an isolated location away from water supplies. Burn bag immediately and stay out of smoke.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-1995

ARC701229
710315MW

STAUFFER CHEMICAL COMPANY

NEW YORK, N.Y. 10017

476-1995

6-2-1971

1/1


NEXT

LABEL

476-1995

6-2-1971

1/1



DYFONATE 10.G

Supplementary Label Claim

476-1995

Peanuts --

For control of lesser cornstalk borer, apply 10 to 14 pounds per acre in a 14-inch band. This is equivalent to 1 to 1.4 pounds of Dyfonate 10.G in 3,730 feet of band 14 inches wide, regardless of row spacing. Apply just prior to pegging and incorporate lightly into the soil.

OR

For control of rootworms and lesser cornstalk borer apply 20 pounds per acre in an 18-inch band over the peanut rows 36 inches apart. Apply just prior to pegging and incorporate lightly into the soil.

NOTE TO REVIEWER: Claim for rootworm control is presently registered.

USDA Reg. No. 476-1995

ARC 710413

80-55