

50 LB.



426-1997 NET
RECEIVED
JUN 13 1969
U.S. FEDERAL BUREAU OF INVESTIGATION
AND INTERNATIONAL TRADE COMMISSION
ECONOMIC POISON REGISTER
REGISTERED UNDER NO. 5562-01
10 ATTACHED CONTROLS

COPPER-SEVIN*-SULFUR 4-10-70 DUST

AN AGRICULTURAL INSECTICIDE & FUNGICIDE

ACTIVE INGREDIENTS:

Copper expressed as metallic (from 5.33% Copper Oxide)	4%
Carbaryl (1-naphthyl N-methylcarbamate)	10%
Sulfur	70%

INERT INGREDIENTS

16%

*Trade-mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate (Carbaryl) 100%

CAUTION **Keep Out of Reach of Children**

HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dusts! Do Not Take Internally!

SKIN CONTACT MAY BE HARMFUL. Avoid contact. Wash hands before eating. Take shower or bath daily after work. Wear regular long-sleeved work clothes. Change to clean clothes daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate irritant to mucous membranes of the throat. Avoid eye contact.

AVOID CONTAMINATION OF FOOD AND FEED. Avoid contact with range of livestock, troughs and watering receptacles.

BURN OR BURY EMPTY CONTAINER — KEEP OUT OF SMOKE OR FUMES.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not use if wind is blowing from directions of travel of spray. Do not use in areas where it is likely to drift to adjacent crops. Do not use in areas where it is likely to drift to adjacent bodies of water. Do not use in areas where it is likely to drift to adjacent residential areas. Do not use in areas where it is likely to drift to adjacent public places. Do not use in areas where it is likely to drift to adjacent schools. Do not use in areas where it is likely to drift to adjacent churches. Do not use in areas where it is likely to drift to adjacent parks. Do not use in areas where it is likely to drift to adjacent playgrounds. Do not use in areas where it is likely to drift to adjacent swimming pools. Do not use in areas where it is likely to drift to adjacent tennis courts. Do not use in areas where it is likely to drift to adjacent golf courses. Do not use in areas where it is likely to drift to adjacent airports. Do not use in areas where it is likely to drift to adjacent highways. Do not use in areas where it is likely to drift to adjacent bridges. Do not use in areas where it is likely to drift to adjacent tunnels. Do not use in areas where it is likely to drift to adjacent overpasses. Do not use in areas where it is likely to drift to adjacent underpasses. Do not use in areas where it is likely to drift to adjacent power lines. Do not use in areas where it is likely to drift to adjacent communication lines. Do not use in areas where it is likely to drift to adjacent utility lines. Do not use in areas where it is likely to drift to adjacent structures. Do not use in areas where it is likely to drift to adjacent vehicles. Do not use in areas where it is likely to drift to adjacent animals. Do not use in areas where it is likely to drift to adjacent humans. Do not use in areas where it is likely to drift to adjacent children. Do not use in areas where it is likely to drift to adjacent pregnant women. Do not use in areas where it is likely to drift to adjacent nursing infants. Do not use in areas where it is likely to drift to adjacent children under 12 years of age. Do not use in areas where it is likely to drift to adjacent children under 6 years of age. Do not use in areas where it is likely to drift to adjacent children under 2 years of age. Do not use in areas where it is likely to drift to adjacent children under 1 year of age. Do not use in areas where it is likely to drift to adjacent children under 6 months of age. Do not use in areas where it is likely to drift to adjacent children under 3 months of age. Do not use in areas where it is likely to drift to adjacent children under 1 month of age. Do not use in areas where it is likely to drift to adjacent children under 1 week of age. Do not use in areas where it is likely to drift to adjacent children under 1 day of age. Do not use in areas where it is likely to drift to adjacent children under 1 hour of age. Do not use in areas where it is likely to drift to adjacent children under 1 minute of age. Do not use in areas where it is likely to drift to adjacent children under 1 second of age. Do not use in areas where it is likely to drift to adjacent children under 1 millisecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 microsecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 nanosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 picosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 femtosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 attosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 zeptosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 yoctosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 ryoctosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 septosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 octosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 nonosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 decasecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 centisecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 millisecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 microsecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 nanosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 picosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 femtosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 attosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 zeptosecond of age. Do not use in areas where it is likely to drift to adjacent children under 1 yoctosecond of age.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment, aircraft or hand duster. Apply at the first sign of infestation and infectio and repeat at 5 to 10-day intervals when needed to maintain control. For the control of diseases persisting after the prescribed time limits have expired, an effective fungicide should be used alone in a manner which will not leave residues in excess of established tolerances.

RECOMMENDATIONS

BEANS: Armyworms, Bea Leaf Beetle, Corn Earworm, Cucumber Beetle, Flea Beetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Velvet Bean Caterpillar — Use 10 to 15 lb. per acre. Mexican Bean Beetle — Use 8 lb. per acre. Apply thoroughly to upper and lower leaf surfaces.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Flea Beetles, Harlequin Bug — Use 8 to 10 lb. per acre. Armyworms, Cabbage Caterpillars, Corn Earworm — Use 10 to 20 lb. per acre. Where cabbage looper is the principal offender, use an alternative pesticide. For control of small loopers after the heads begin to form, apply every 7 to 7 days. **Do not apply within 3 days of harvest.**

CORN: Corn Earworm, Corn Rootworm (adults), European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhoppers, Sap Beetles — Use 10 to 20 lb. per acre. For Corn Earworm, apply when silks first appear and repeat at 2 to 3-day intervals throughout the silk-up period. Direct duster application toward ear zone to insure thorough coverage of leaf surface.

COTTON: **Early-Season Control** — Cotton Fleahopper, Cotton Leafworm, Flea Beetles, Striped Blister Beetle, Thrips — Use 10 to 20 lb. per acre. **Mid- and Late-Season Control** — Boll Weevil, Fall Armyworm, Leafhoppers, Leaf Rollers, Tarnished Plant Bug — Use 10 to 20 lb. per acre, depending on size of cotton and plant population. Apply every 7 to 14 days during peak season. Pink Bollworm — Use 10 to 20 lb. per acre every 7 to 7 days, depending on size of cotton and population. Salt-marsh Caterpillar, Stinkbug — Use 20 lb. per acre. **Do not apply within 7 days of grazing or harvesting crop residue for feed or bedding for dairy or meat animals.**

EGGPLANT, PEPPER, POTATO — Colorado Potato Beetle, Flea Beetles, Leafhoppers — Use 10 to 20 lb. per acre. European Corn Borer, Fall Armyworm, Lace Bugs, Stinkbugs, Tarnished Plant Bug, Tomato Fruitworm, Tomato Hornworm — Use 10 to 20 lb. per acre.

PLAUIIS — Bean Leaf Beetle, Blister Beetle, Cucumber Beetle, Japanese Beetle, Leafhoppers, Thrips, Velvet Bean Caterpillar — Use 10 to 20 lb. per acre. Armyworms, Corn Earworm, Grasshoppers, Stinkbugs, Webworms — Use 10 to 20 lb. per acre. Peanut Leaf Spot — Apply dust during emergence after planting or as soon as signs of damage appear. Repeat every 7 to 10 days. Total application should not exceed 30 lb. per acre.

NOTICE: This product is a restricted-use pesticide. It is not to be used on any particular crop, animal, or human without the specific instructions which are contained on Stauffer's label.

STAUFFER CHEMICAL CO. NEW YORK