



# Sutan 10 GRANULAR

50 LB. NET

A SELECTIVE HERBICIDE FOR CORN (Field, Sweet, & Silage)

## A SELECTIVE HERBICIDE

### ACTIVE INGREDIENT:

S-ethyl diisobutylthiocarbamate 10.0%

### INERT INGREDIENTS:

90.0%

Contains Ten Pounds Active Ingredient per 100 Pounds 100%

U. S. Patent No. 2,913,327

## CAUTION

Keep Out of Reach of Children

AVOID INHALATION OF DUST.

HARMFUL IF SWALLOWED.

AVOID CONTACT WITH SKIN, EYES AND CLOTHING

WASH THOROUGHLY AFTER USE.

DO NOT CONTAMINATE FEED AND FOOD.

### GENERAL USE PRECAUTIONS

#### READ ENTIRE LABEL BEFORE USING

Applied according to directions and under normal growing conditions, SUTAN will not harm the treated crop. During germination and early stages of growth, extended periods of unusually cold and wet or hot and dry weather, insect or plant disease attack, the use of certain soil-applied systemic insecticides, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. SUTAN used under these abnormal conditions could result in crop injury.

SUTAN is recommended for use on mineral soils only.

Do not apply in combination with fertilizers, insecticides or fungicides unless the compatibility has been tested and proven to be satisfactory prior to use.

Further information can be obtained from your Stauffer dealer.

Do not store near seeds or fertilizers.

Do not contaminate irrigation water or water used for domestic purposes.

Keep container closed when not in use.

### WEEDS CONTROLLED

#### Annual Grasses:

|                           |                         |
|---------------------------|-------------------------|
| Barnyardgrass; Watergrass | (Echinochloa crusgalli) |
| Crabgrass                 | (Digitaria spp.)        |
| Giant Foxtail             | (Setaria faberii)       |
| Yellow Foxtail            | (Setaria glauca)        |
| Green Foxtail             | (Setaria viridis)       |
| Goosegrass                | (Eleusine indica)       |
| Seedling Johnsongrass     | (Sorghum halepense)     |

#### Perennial Weeds:

|                 |                      |
|-----------------|----------------------|
| Purple Nutgrass | (Cyperus rotundus)   |
| Yellow Nutgrass | (Cyperus esculentus) |

Existing stands of nutgrass must be turned under and chopped up thoroughly before treatment. Prior to planting, SUTAN 10 Granular must be thoroughly mixed into the soil.

476-2001  
ACCEPTED  
May 5 1968  
UNDER THE FEDERAL INSECTICIDE ACT  
FEDERAL BUREAU OF INVESTIGATION  
WASHINGTON, D. C. 20535

### DIRECTIONS FOR USE

SUTAN is a selective herbicide which is mixed (incorporated) into the soil for control of weeds listed on this label. SUTAN controls weeds as their seeds germinate by interfering with normal germination and seedling development. It does not control established weeds. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

### APPLICATION DIRECTIONS

Apply only to well worked soil that is dry enough to permit thorough mixing with incorporation equipment. Apply to a dry soil surface only. Apply uniformly the recommended rate of SUTAN 10 Granular using equipment designed for application of granular herbicides. Equipment should be carefully calibrated before use and checked frequently during application to be sure equipment is working properly and delivering a uniform distribution pattern. Avoid overlaps that will increase SUTAN dosage above recommended rates as plant injury may occur.

In irrigated areas, do not apply SUTAN before pre-irrigation.

### INCORPORATION DIRECTIONS

SUTAN 10 Granular must be incorporated into the soil immediately to prevent loss of the herbicide. Whenever possible, application and incorporation should be done in the same operation.

**SOIL MIXING (INCORPORATION) BEFORE PLANTING** Use power-driven cultivation equipment set to cut to a depth of three inches; OR a tandem disc set to cut to a depth of six inches followed by a spiked-tooth harrow or some other leveling device which extends beyond the ends of the disc. For thorough mixing, disc in two different directions (cross disc).

### CULTURAL PRACTICES FOLLOWING APPLICATION

SUTAN is not persistent in the soil and susceptible weeds germinating late in the season may not be controlled. Cultivate to control these late germinating susceptible weeds, as well as those weeds not susceptible to SUTAN. Do not cultivate deeper than depth of herbicide incorporation.

### RECOMMENDATIONS

**CORN:** (Field, Sweet & Silage) Apply and incorporate SUTAN just before planting (pre plant). Plant seed to a maximum depth of two inches. Do not use on milo or sorghum or hybrid corn grown for seed.

**Broadcast rate per acre:** For control of annual grasses and nutgrass, apply SUTAN 10 Granular at 40 pounds per acre. One cultivation should be made when the corn plants are three to six inches tall.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

U.S.D.A. Reg. No. 476-2001

ARC 680209

STAUFFER CHEMICAL COMPANY  
NEW YORK, NEW YORK

87 O Ag 680325

BURN EMPTY BAG • KEEP OUT OF SMOKE

BEST DOCUMENT AVAILABLE