

Net Contents

(STAUFFER SHIELD)

P R E F A R [®] 4E

A Selective Pre-plant Herbicide for Control of Annual Grasses and Broadleaf Weeds

Contains 4 Pounds PREFAR per Gallon

ACTIVE INGREDIENT:

S-(O,O-diisopropyl phosphorodithioate) of N-(2-mercaptoethyl) benzenesulfonamide 45.2%

INERT INGREDIENTS 54.8%

100.0%

U. S. Patent Number 3,205,253

C A U T I O N


KEEP OUT OF REACH OF CHILDREN

4/24/68 MSZ Harmful if swallowed. Avoid ~~prolonged or repeated~~ contact with the skin. Avoid contact with the eyes. Avoid contamination of food and feed, drinking water, fish ponds, et cetera. Wash thoroughly after use.

Do not use or store near heat or open flame.

Minimum storage temperature 42°F.*

* Product crystallizes at lower temperatures. Warm or store at higher temperatures and mix to redissolve crystals and assure uniformity before use.



STAUFFER CHEMICAL COMPANY

New York

WEEDS CONTROLLED

Crabgrass	(<u>Digitaria</u> spp.)
Foxtail	(<u>Setaria</u> spp.)
Jungle Rice	(<u>Echinochloa</u> <u>colonum</u>)
Watergrass	(<u>Echinochloa</u> <u>crusgalli</u>)
Lambsquarters	(<u>Chenopodium</u> <u>album</u>)
Pigweed	(<u>Amaranthus</u> spp.)
Purslane	(<u>Portulaca</u> <u>oleracea</u>)

GENERAL USE PRECAUTIONS

READ ALL LABEL DIRECTIONS BEFORE USING.

PREFAR should be used only for recommended purposes and at recommended rates. (DO NOT OVERDOSE)

Do not store near seeds or fertilizers.

Keep container closed when not in use.

Use on mineral soil only. PREFAR MUST BE THOROUGHLY MIXED INTO THE SOIL.

Do not contaminate irrigation water or water used for domestic purposes.

*4/27/68
MSL* ~~Fish may be killed by contamination of streams or ponds with PREFAR. This product is toxic to fish. Keep out of lakes, ponds, streams, tidal marshes and estuaries.~~

All crops specified on the label may be planted in rotation but do not apply PREFAR 4E more often than once every 12 months. To avoid injury to crops not specified on the label do not plant until 18 months after the last PREFAR 4E application.

DIRECTIONS FOR USE

SOIL PREPARATION: Prepare the soil for seeding according to good agricultural practice.

APPLICATION AND SOIL INCORPORATION: PREFAR must be thoroughly incorporated to a depth of one to two inches before planting. Spray equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and are delivering a uniform spray pattern. Apply uniformly the recommended rate of PREFAR in 20-100 gallons of water per acre. Apply to well-worked soil that is dry enough to permit thorough incorporation.

RECOMMENDATIONS

Crop	When to Apply	Dosage - Quarts/ Broadcast Acre	Remarks
LETTUCE (Head or Leaf)	Pre-plant	5-6	For control of all weeds listed. Apply and incorporate on bed tops with power-driven tillers.
CUCUMBERS SUMMER SQUASH CANTALOUPE MUSKMELONS WATERMELONS CRENSHAW MELONS PERSIAN MELONS	Pre-plant	5-6	For control of all weeds listed. On bedded crops apply and incorporate on bed top with power-driven tillers. OR On flat planted crops, apply (in bands or broadcast) and then incorporate with equipment which will result in thorough incorporation.
COTTON	Pre-plant	2	For control of annual grasses in Southwestern States. Use only on light sandy soils. Apply to the flat soil surface before furrowing (listing) for the pre-planting irrigation and then continue with normal practices. OR Apply after furrowing (listing) and before the pre-planting irrigation. Incorporate with a sectioned, rolling cultivator. OR Apply after the pre-planting irrigation and before the final seed bed preparation.

STANDARD STAUFFER DISCLAIMER