POISON

IF SWALLOWED [illegible] glass of warm water [illegible] repeat [illegible] have victim lie down and [illegible]

INHALATION [illegible] from contaminated atmosphere.

SKIN [illegible] of contact, remove contaminated clothing and [illegible] wash skin thoroughly with soap and water.

EYES [illegible] immediately with water for at [illegible]

INFORMATION FOR PHYSICIAN:

[illegible] maintain [illegible] of secretions, maintenance of a [illegible] airway and [illegible] artificial respiration. **When cyanosis has been relieved**, atropine may be administered in large [illegible] repeated as necessary to the point of tolerance [illegible] symptoms warrant further treatment. Protopam chloride (2-PAM) [illegible] **has shown utility as adjunctive therapy**. [illegible] may [illegible] delayed absorption of the poison [illegible] after initial [illegible] have been reported after initial [illegible] every [illegible] examination **of the patient is indicated** [illegible] at least 48 to 72 hours.

WARNING

[illegible] protective clothing and goggles [illegible] of a type passed by the U.S. Department [illegible] parathion protection. Airplane pilots [illegible] self-contained mask.

[illegible] portion [illegible] of operating areas or vicinity [illegible] border of crops. Vegetated areas should not be [illegible] insecticide and volatile pesticides have [illegible] fruit trees in dense orchards until [illegible] been allowed for dissipation of vapors [illegible]

[illegible] with soap and water before [illegible] Wash [illegible] contaminated clothing with [illegible]

LOT [illegible]

OBSERVE THE FOLLOWING USE PRECAUTIONS

[illegible] use limitations given in the label for specific crops [illegible] in order that pesticide residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration. Use only at recommended rates and intervals, and [illegible] not apply closer to harvest than [illegible] specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for **damages caused** by drift from his property to that of others.

[illegible] this product on feed or forage to be fed to dairy animals or animals being finished for slaughter.

This product is toxic to fish and wildlife. Keep out lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated.

To protect bees, do not apply when crops are in bloom. Remove or cover hives if application is necessary during bloom.

For best results apply when foliage is dry and there is little or no wind.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as to the timing, number and rate of applications needed will vary with local [illegible]

DESTROY EMPTY CONTAINER — NEVER RE-USE

Completely empty contents and bury unused chemical 18 inches deep in an isolated location away from water supplies.

GLASS CONTAINER: Break container and bury 18 inches deep.

METAL CONTAINERS: 1-Gal Can — Pour 1 qt. of water into empty can. Add 1 tablespoon of household detergent. Rotate can carefully until all inner surfaces are wet. Bury rinse solution 18 inches deep. Punch holes in top and bottom of container, crush and bury. **5-Gal. Drum** — Pour 2 qt. of water into empty container. Slowly add ½ cup caustic soda (lye) and 2 tablespoons household detergent. Then proceed as for 1-gal. container. **30-Gal. Drum** — Same as for 5-gal. drum except use 3 gal. water, 1 lb. lye, and ½ cup detergent for rinsing. **55-Gal. Drum** — Same as for 5 gal. drum except use 5 gal. water, 2 lb. lye, and 1 cup detergent for rinsing.

Do not get rinse solution on [illegible] clothing, or in eyes. In case of contact, immediately wash with soap and water.

6-25-68
476-2015

Stauffer CHEMICALS

PARATHION-THIODAN 1-2 E

EMULSIFIABLE LIQUID — AGRICULTURAL INSECTICIDE

(Contains 1 lb. Parathion and 2 lb. Thiodan per gallon)

ACTIVE INGREDIENTS:	
Parathion (O, O-Diethyl O-p-nitrophenyl phosphorothioate)	11.5%
Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	22.8%
Xylene Range Aromatic Solvent	60.0%
INERT INGREDIENTS	5.7%
	100.0%

Thiodan is a Registered Trademark of Canadian Hoechst Ltd. U.S. Patent No. 2,799,685

POISON

POISON — Keep Out of Reach of Children

DANGER

POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN OR EYES. RAPIDLY ABSORBED THROUGH SKIN OR EYES. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING
DO NOT BREATHE VAPOR OR SPRAY MIST
POISONING SYMPTOMS: [illegible]
ANTIDOTE: Atropine [illegible]
CALL A PHYSICIAN IMMEDIATELY [illegible]
Parathion is a highly toxic cholinesterase inhibitor.
Thiodan [illegible]
BEFORE USING SEE FIRST AID TREATMENT ON LEFT PANEL OF THIS LABEL.

PROTECT FROM TEMPERATURES BELOW 25°F.

NOTE BEFORE USING, AGITATE CONTENTS BY SHAKING OR ROLLING CONTAINER.

*Product strength [illegible] guaranteed [illegible] to be useful condition [illegible]

STAUFFER CHEMICAL CO., NEW YORK

[illegible] GALLONS NET

Stauffer CHEMICALS

PARATHION-THIODAN 1-2 E

EMULSIFIABLE LIQUID — AGRICULTURAL INSECTICIDE

(Contains 1 lb. Parathion and 2 lb. Thiodan per gallon)

ACTIVE INGREDIENTS:	
Parathion (O, O-Diethyl O-p-nitropheny phosphorothioate	11.5%
Endosulfan (Hexachlorohexahydromethan >2,4,3-benzodioxathiepin oxide)	22.8%
Xylene Range Aromatic Solvent	60.0%
INERT INGREDIENTS	5.7%
	100%

Thiodan is a Registered Trademark of Canadian Hoechst, Ltd. U.S. Patent No. 2,799,685

POISON **Keep Out of Reach of Children**

DANGER

POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN OR EYES. RAPIDLY ABSORBED THROUGH SKIN OR EYES. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING
DO NOT BREATHE VAPOR OR SPRAY MIST

POISONING SYMPTOMS: Headache, [illegible] the chest, vomiting, [illegible]

ANTIDOTE: Atropine is the specific antidote. Obtain a hypodermic syringe of Atropine Sulfate tablets from your physician. Keep it ready.

CALL A PHYSICIAN IMMEDIATELY • person handling [illegible] exposed. Early administration of Atropine is important. Do not wait for symptoms.

Parathion is a highly toxic cholinesterase inhibitor.
Thiodan produces [illegible]

BEFORE USING SEE FIRST AID TREATMENT ON LEFT PANEL OF THIS LABEL

PROTECT FROM TEMPERATURES BELOW 28° F.*

NOTE: BEFORE USING, AGITATE CONTENTS BY SHAKING OR ROLLING CONTAINER.

*Product freezes but is not damaged by low temperatures and will regain to useful condition without application of heat or thawing out from storage at 0° F.

STAUFFER CHEMICAL CO., NEW YORK

_____ GALLONS NET

D[illegible]

A[illegible] mixtures [illegible]

For [illegible] name [illegible] directions [illegible] recommendations [illegible] or [illegible] consult [illegible] product [illegible] package.

RECOMMENDATIONS

Dosage [illegible] athlo[illegible] rates: [illegible] gallons per acre.

BROCCOLI, CABBAGE [illegible] Cabbage Loopers, Cross-striped Cabbageworm, Diamond-back Moth Larvae, Imported Cabbageworm, Harlequin Bug, Imported Cabbageworm, Striped Blister Beetle. Use 1½ qts. per acre. Do not apply within 7 days of harvesting Broccoli, Brussels Sprouts or Cabbage.

CAULIFLOWER Cabbage Loopers, Cross-striped Cabbageworm, Diamond-back Moth Larvae, Imported Cabbageworm, Harlequin Bug, Striped Blister Beetle. Use 1½ qts. per acre. Do not apply within 7 days of harvesting Cauliflower.

CUCUMBERS, PUMPKINS, SUMMER SQUASH, WINTER SQUASH Aphids, Cucumber Beetles, [illegible] Use 1½ qts. per acre. Do not apply before vining. Do not [illegible]

HEAD LETTUCE: Cabbage Loopers, Cross-striped Cabbageworm, Green Peach Aphids, Potato Flea Hopper and Cabbage Looper. Use 1½ qts. per acre. Do not apply within 3 days of harvesting Head Lettuce. Remove wrapper leaves at harvest. Do not feed treated lettuce to livestock. Do not [illegible]

MELONS: [illegible] Cabbage Loopers, [illegible] Do not apply [illegible]

Cr[illegible] and [illegible] are [illegible]

TOBACCO [illegible] Cutworms, Hornworms, Flea Beetles, Grasshoppers, Leaf Eating Caterpillars, Leaf Miners, Thrips, Wireworms. Use 1½ qts. per acre. Apply as a dilute spray to 2 qts. per acre. Apply as a dilute spray. Use 2 qts. per acre.

NOTE: Do not [illegible] within 5 days of [illegible] or harvest. Do not graze [illegible] within 24 hours after application of this product. Do not [illegible] following the application of this product.

TOBACCO (PLANT BED TREATMENT. [illegible] 1 pt. per 100 square feet. Apply as a drench using 1 gallon of spray per 100 square feet of plant bed.

NOTICE: [illegible] or implied, [illegible] of any person [illegible] these which are [illegible]

U.S. Patent No. 2,799,685

Made in U.S.A.
Stauffer Chemical Co., New York