

476
WIDE RANGE
ECONOMIC FUNGICIDE

PRECAUTIONS

DIRECTIONS FOR USE

RECOMMENDATIONS

CROP USES

TREE FRUIT CROPS

APPLES
APRICOTS
CHERRY
ORANGE
PEACH
PLUM
PRAIRIE
RAISIN
STRAWBERRY

GRAPES
LEMON
LIME
PEACHES
PEARS
PINEAPPLE

VEGETABLE AND FIELD CROPS
ALFALFA
BEANS
CUCUMBER
EGGPLANT
MELON
PUMPKIN
SQUASH
TOMATO

GRASS AND CEREAL CROPS
Barley, corn, oats, wheat, barn grass,
canary grass, clover, cordgrass, red top, timothy, and yellow
foxtail

PEPPERS
POTATO
SWEET POTATOE
WHEAT

FRUIT
GROUNDCOVER
ORCHARD FLOOR
SUGAR BEET
TANSEY

FRUIT
TANSEY

FRUIT
TANSEY

CORN
COTTON

CRANBERRIES
CUCUMBER
PUMPKIN
SQUASH

EGGPLANT
GRASS AND CEREAL CROPS

PEPPERS
POTATO

FRUIT
TANSEY

FRUIT
TANSEY

FRUIT
TANSEY

FRUIT
TANSEY

CYT

THE PREMIUM AGRICULTURAL EMULSIFIABLE

ACTIVE INGREDIENTS:
Malathion, O-dimethyl S of diethyl mercaptosuccinylsulfonamide Aromatic S
INERT INGREDIENTS:
CONTAINS 5 LBS. PREMIUM GRADE MALATHION PER GAL

CAUTION

HARMFUL TO WATER LIVING ORGANISMS
Wash thoroughly after handling
Do not eat, drink, or smoke while using this product
Do not breathe vapors
Do not get on clothing
Do not use near water

NOTICE: Do not use this product on crops, plants, or animals that are being raised for human consumption. Do not use on crops, plants, or animals that are being raised for human consumption.

STUFFE



CYTHION^(R) 5-E

THE PREMIUM GRADE MALATHION^(R) AGRICULTURAL INSECTICIDE EMULSIFIABLE LIQUID

CONTAINS
CYTHION

ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl phosphorothioate) of diethyl mercaptopyruvate
Xylene Range Solvent

INERT INGREDIENTS:

CONTAINS 5 LBS. PREMIUM GRADE MALATHION PER GAL.

Do not use, pour, spill or store near heat or open flame.

Rinse spray equipment and containers. Discharge liquid waste in pit in non-crop lands located away from water supplies.

PERFORATE EMPTY METAL CONTAINERS CRUSH & BURY GLASS.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Avoid contact with vapor or spray mist. Avoid contact with skin. Wash thoroughly with soap and water before eating, drinking, or using tobacco. Do not contaminate food and feed.

NOTE TO PHYSICIANS:

For a list of physicians in your area, contact your local Stauffer distributor.

Keep Out of Reach of Children

Store in original container. Do not use if container is damaged. Do not use if contents are not as labeled. Do not use if contents are not as labeled.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranty of merchantability or of fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Lot No.

1 GAL. NET

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

STRAWBERRIES

Apply to plants in fruiting stage. Spray thoroughly. Repeat every 7-10 days. California only.

SWISS CHARD

Apply to plants in fruiting stage. Spray thoroughly. Repeat every 7-10 days. California only.

TOMATOES

Apply to plants in fruiting stage. Spray thoroughly. Repeat every 7-10 days. California only.

PASTURE AND RANGE LAND

Apply to pastures and range lands. Not grazed. California only.

WASTELANDS, ROADSIDES, SOIL-BANK LANDS Not grazed. California only.

GRAINS: BARLEY, CORN, GRAIN, SOYBEAN SEED, OATS, RICE, RYE, WHEAT. RESIDUAL SPRAY. BEFORE STORAGE.

GRAINS: GOING INTO STORAGE. AFTER STORED.

RESIDUAL SPRAY FOR BORDERS OF BAG FLOUR AND PACKAGED CEREALS.

FIELD AND GARDEN

Apply to plants in fruiting stage. Spray thoroughly. Repeat every 7-10 days. California only.

PEANUTS Apply to plants in fruiting stage. Spray thoroughly. Repeat every 7-10 days. California only.

PEANUTS BULK SPRAY TREATMENT.

PEANUTS RESIDUAL SPRAY.

PEANUTS PREVENTATIVE SURFACE TREATMENT.

PEANUTS RESIDUAL SPRAY.

PEANUTS RESIDUAL SPRAY.

PEANUTS RESIDUAL SPRAY.

PEANUTS RESIDUAL SPRAY.

PEANUTS RESIDUAL SPRAY.

PEANUTS RESIDUAL SPRAY.

STORAGE PEST PROTECTION

ORNAMENTALS

FIELD AND GARDEN SEEDS

FIELD AND GARDEN SEEDS

PEANUTS

PEANUTS RESIDUAL SPRAY BEFORE STORAGE

PEANUTS BULK THERAY TREATMENT GOING INTO STORAGE

PEANUTS SURFACE TREATMENT SURFACE SPRAY

ORNAMENTALS

Decorative are use in 100 gal. of water spray

POULTRY

POULTRY RANGES Apply 100% Make full coverage

ROOST PAINT Apply 100% Make full coverage

FLY CONTROL

Flies in and around facilities which house domestic animals...

STRAIGHT CYTHIUM SPRAYS WITH ON BAIT SPRAYS

Concentration	Application	Remarks
100%	Direct spray	For immediate control
100%	On bait	For long-term control

CAUTION

GRUBS, FLIES AND DRIED FRUIT BEETLES

MOSQUITO CONTROL

MOSQUITOES, FLIES AND SMALL FLYING INSECTS

MOSQUITO LARVAE IN STANDING WATER