

(STANDARD 300-1)

EPTAM[®]

1,1-DIMETHYL-2,2-DICHLOROETHAN-1-OL

USE ONLY IN ACCORDANCE WITH DIRECTIONS ON LABEL. DO NOT USE FOR CONTROL OF ANIMALS OR FOR ANY OTHER PURPOSES NOT SPECIFIED ON LABEL.

CONTAINS 6 FLUID OUNCES (170.1 ML) OF EPTAM PER GALLON.

ANALYTICAL DATA:

SUBSTANCE IDENTIFICATION: 76.9%
INERT INGREDIENTS: 23.0%
100.0%

U. S. PATENT NO. 2,703,217

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED
AVOID CONTACT WITH SKIN AND CLOTHING
WASH THOROUGHLY WITH SOAP
AVOID INHALATION OF FOG
DO NOT CONTAMINATE FOOD OR FEED

CAUTION

DO NOT USE IN OPEN

DO NOT USE

DO NOT USE IN OPEN
DO NOT USE IN OPEN
DO NOT USE IN OPEN

EPTAM 6-E (ORGANIC)

2.

OBSERVE THE FOLLOWING PRECAUTIONS:

READ ALL LABEL DIRECTIONS BEFORE USING.

EPTAM IS TO BE USED ONLY FOR THE PURPOSES SPECIFIED ON LABEL. DO NOT USE AT RATES OTHER THAN THOSE SPECIFIED.

EPTAM IS RECOMMENDED FOR USE ON MINERAL SOILS ONLY.

FOR ALL USE, EPTAM MUST BE THOROUGHLY MIXED INTO THE SOIL.

EPTAM MAY CAUSE INJURY TO CRUSTACEANS UNDER CERTAIN SOIL AND CLIMATIC CONDITIONS OR IF DIRECTIONS ARE NOT FOLLOWED.

KEEP CONTAINER CLOSED WHEN NOT IN USE. DO NOT STORE NEAR SEEDS OR FERTILIZERS.

RINSE SPRAY EQUIPMENT AND CONTAINER. BURY LIQUID WASTE IN PIT IN NON-CROP LANDS LOCATED AWAY FROM WATER SUPPLIES.

DO NOT RE-USE EMPTY CONTAINER. PERFORATE EMPTY METAL CONTAINERS.

CRUSH AND BURY CLIPS IN SAME PLACE.

EPTAM CAN BE USED ON THE FOLLOWING PLANTS:

HERBACEOUS PLANTS

ALYSSUM	BALSAM	DAYLILY	PANSY
AGERATUM	BECONIA	DIANTHUS	PELARGONIUM
AMARANTHUS	CHRYSANTHEMUM	NARCISSUS	ZINNIA
ASTERS	DANIELA	MADEIRA	

GRASSY PLANTS

AJUCA	EAGLE
GAZANIA	PERILLA
HYPERICUM	SEDUM
ICE PLANT	STREPTOCARPUS
IVY	

EVERGREEN PLANTS

BLUEBERRY	ELDERBERRY	FRUIT	FRUIT
DOGWOOD	FIR	FRUIT	FRUIT
GERANIUM	FRUIT	FRUIT	FRUIT
GRASS	FRUIT	FRUIT	FRUIT
HEARTSEASE	FRUIT	FRUIT	FRUIT
HEARTSEASE	FRUIT	FRUIT	FRUIT
HEARTSEASE	FRUIT	FRUIT	FRUIT
HEARTSEASE	FRUIT	FRUIT	FRUIT

DO NOT USE IN OPEN
DO NOT USE IN OPEN
DO NOT USE IN OPEN

ACCEPTED

HOW TO USE

PRE-PLANT APPLICATION - ROTOTILLER.

POST-PLANT APPLICATION - FOR ANNUAL WEEDS USE HAND RAKE OR HOE OF WATER IN IMMEDIATELY AFTER APPLICATION TO A DEPTH OF 2 TO 3 INCHES. WHERE INCORPORATION BY HAND RAKING IS DONE, LIGHT WATERING AFTER RAKING IS RECOMMENDED.

FOR PERENNIAL WEEDS, INCORPORATE TO A DEPTH OF 6 INCHES WITH A ROTOTILLER.

RECOMMENDATIONS

FOR ANNUAL WEED CONTROL - USE EPTAM 6-E AT THE RATE OF $3\frac{1}{3}$ QUARTS IN 30 TO 60 GALLONS OF WATER PER ACRE ($\frac{1}{3}$ CUP ($2\frac{2}{3}$ FLUID OUNCES) PER 1,000 SQUARE FEET).

FOR QUACKGRASS, HULMGRASS, AND PERENNIALGRASS CONTROL IN TREES AND SHRUBS ONLY - EXISTING STANDS OF THESE PERENNIAL GRASSES MUST BE TURNED UNDER AND CHOPPED UP THOROUGHLY BEFORE TREATMENT. USE EPTAM 6-E AT THE RATE OF 1 GALLON IN 30 TO 60 GALLONS OF WATER PER ACRE ($\frac{3}{8}$ CUP (3 FLUID OUNCES) PER 1,000 SQUARE FEET).

FOR MUGWORT (GERYSALETHOIFERED) CONTROL IN THE FOLLOWING PLANTS: JUNIPER, JAPANESE LILLY, LILY, PEARLY LILY, PINK LILY - USE $2\frac{1}{2}$ GALLONS OF EPTAM 6-E IN 30 TO 60 GALLONS OF WATER PER ACRE ($\frac{7}{8}$ CUP (7 FLUID OUNCES) PER 1,000 SQUARE FEET). MIX THOROUGHLY INTO THE TOP 6 INCHES OF SOIL. APPLY 4 WEEKS BEFORE DESIRED PLANTING DATE.

WHEN TO USE EPTAM

HERBACEOUS PLANTS AND CROPPED CORN: APPLY 2 WEEKS BEFORE TRAIL - PLANTING OR AFTER CROPPED STARTS IN THE SPRING.

TREES AND SHRUBS: APPLY 2 WEEKS BEFORE TRAIL PLANTING DATE AND CROPPED STOCK (ONLY) AND ANYTIME AFTER TRAIL PLANTING. ALSO, ESTABLISHED PLANTS: APPLY AFTER CROPPED STARTS IN THE SPRING.

NOTE: STRENGTH OF EPTAM 6-E MAY VARY IN DIFFERENT FORMULATIONS. THEREFORE, THE USER SHOULD CONSULT THE LABEL FOR ANY PARTIAL RATE. EPTAM 6-E IS A TRADE MARK OF EPTAM, INC. WHICH IS CONTAINED IN STAMPS OF 17.11.

ACC. ED.