




# CAPTAN-LINDANE 12.5-25

## SEED PROTECTANT

FUNGICIDE — INSECTICIDE — DRY TREATMENT APPLICATION

SPECIAL LUBRICANT ADDED — DISCOLORED



### ACTIVE INGREDIENTS:

Captan: N-[[trichloromethyl]thio]-4-cyclohexene-1,2-dicarboximide*	12.5%
Gamma isomer of Benzene Hexachloride (from Lindane)	25.0%

### INERT INGREDIENTS:

	62.5%
	100.0%

\*Not licensed for use in countries where valid patents still exist.

### WARNING

Keep Out of Reach of Children

MAY BE FATAL IF SWALLOWED.

MAY BE ABSORBED THROUGH THE SKIN.

Do not breathe dust.

Apply dust out-of-doors or in a well ventilated room.

Avoid contact with skin and eyes.

Wash thoroughly after handling.

Do not contaminate food or feed.

**NOT FOR USE OR STORAGE IN OR AROUND THE HOME.**

**BURY WASTE AND EMPTY CONTAINER IN AN ISOLATED AREA AWAY FROM WATER SUPPLIES.**

*NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.*

2-ARC 72/31 35

EPA Reg. No. 476-2036

1 LB. NET

### OBSERVE THE FOLLOWING USE PRECAUTIONS

Read all label directions before using and use only as specified on this label.

Dispose of excess treated seed by burial.

**FISH AND WILDLIFE CAUTIONS:** This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Birds feeding on treated seeds may be killed.

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED.

DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.

The federal government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed. Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

### DIRECTIONS FOR USE

This formulation is for use in the dry powder form for dust treatment of seeds for protection against seed rot, damping-off, seed corn beetles, seed corn maggots and wireworms. Mix seed with recommended amount of this material, in any commercial seed treader or barrel treader, until thoroughly coated. DO NOT MIX WITH HANDS. May be used on seed previously treated with a fungicide. Avoid lengthy storage of treated seed to prevent loss of LINDANE. If soil temperature is extremely high due to low moisture, hot weather or mid summer planting time, reduce the following dosage rates by one-half:

#### AMOUNT OF CAPTAN-LINDANE 12.5-25 SEED PROTECTANT TO USE ON SEEDS

✓ Barley	4 oz. per 100 lb.
✓ Corn	4½ oz. per bu.
✓ Cotton	3 oz. per 100 lb.
✓ Safflower	2 oz. per 100 lb.
✓ Sunflower	4 oz. per 100 lb.
✓ Sugar Beets	16 oz. (1 lb.) per 100 lb.

Made in U.S.A. by

# STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880