



**ACCEPTED**  
 MAY 28 1970  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTER-  
 ED UNDER NO. 476-2038

# WETTABLE SULFUR

## AGRICULTURAL INSECTICIDE-FUNGICIDE

WETTABLE SULFUR — FOR MIXING WITH WATER

**ACTIVE INGREDIENT:**

Sulfur

97.0%

**INERT INGREDIENTS:**

3.0%

100.0%

### CAUTION

Keep Out of Reach of Children

### USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on apricots, Anjou and comice pears, cranberries, cucurbits (cucumbers, squash and some varieties of melons), filberts, spinach, tung trees, walnuts and other sensitive plants.

Do not apply when shade temperatures exceed or are likely to exceed 90° F.

Sulfur may burn foliage and fruit during periods of high temperatures and under certain climatic conditions.

Do not use sulfur within 30 days of an oil spray, 60 days on citrus.

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

### DIRECTIONS FOR USE

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank, or pre-mix in a bucket half filled with water and pour mixture into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, and spraying operations. Do not allow mixture to stand. Do not use in low-volume gear-type spray equipment. Do not combine with emulsifiable liquids in the same spray tank.

### RECOMMENDATIONS

**APPLES:** Scab, powdery mildew. Apply 4-5 lbs. per 100 gals. from pink or preblossom spray through second cover spray. Add 1 gal. lime-sulfur in pink and calyx sprays. Use on sulfur-tolerant varieties only.

**CHERRY:** Blossom blight, brown rot, leaf spot, powdery mildew. Apply 4-6 lbs. per 100 gals. starting at shuck-fall.

**CITRUS:** Six-spotted and citrus rust mites. Apply 4 lbs. per 100 gals. citrus flat and Yuma spider mites. Apply 4 lbs. per 100 gals. (At least 60 lbs. per acre.)

**GRAPES:** Powdery mildew (eradicator spray). Apply 2 lbs. per 100 gals. plus a wetting agent. Care should be taken to wet all surfaces from which the fungus is to be eradicated.

**PEAR:** Scab, powdery mildew. Apply 4 lbs. plus 1 gal. lime-sulfur per 100 gals. in the pink or pre-blossom spray. On Bartlett pears, apply 4-6 lbs. at petal fall through second cover sprays.

**PEACH, NECTARINE:** Brown rot, botryneum blight, rust, peach silver mite, powdery mildew. Apply 5-6 lbs. per 100 gals. from popcorn stage up to harvest. For brown rot control in summer sprays apply up to 10 lbs. per 100 gals. starting about 4-5 weeks before harvest and continuing at weekly intervals.

**PLUM, PRUNE:** Brown rot, leaf spot, prune rust. Apply 5-6 lbs. per 100 gals. in summer sprays starting about 5 weeks before harvest.

**STRAWBERRIES:** Powdery mildew. Apply 1-5 lbs. per 100 gals. Discontinue application during hot weather.

### ORNAMENTALS

**ROSES, PERENNIALS, SHRUBS:** Black spot, powdery mildew, rust. Apply 3 tablespoons per gallon (6 lbs. per 100 gals.) and spray for full coverage wetting both sides of leaves. Begin applications when new growth appears in the spring and repeat every 7 days, if necessary.  
**CAUTION:** Some burning may occur when used during warm weather.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-2038

50 LB. NET

NET WT. 50 LB.

**STAUFFER CHEMICAL COMPANY** NEW YORK, NY 10017



ACCEPTED  
MAY 28 1971  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROTENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 172-2038

# WETTABLE SULFUR

## AGRICULTURAL INSECTICIDE-FUNGICIDE

WETTABLE SULFUR — FOR MIXING WITH WATER

ACTIVE INGREDIENT:	
Sulfur	97.0%
INERT INGREDIENTS:	3.0%
	100.0%

### CAUTION

Keep Out of Reach of Children

#### USE PRECAUTIONS

Sulfur will cause severe frost and leaf injury to sulfur-sensitive plants. Do not use on sensitive plants. Apply and cultivate fruits, ornamentals, rabbits, guinea pigs, quail, and some varieties of melons, peaches, pears, apples, walnuts, and other nuts to the plant.

Do not apply when wind velocity is greater than 10 miles per hour.

Sulfur may cause frost and fruit damage on plants at low temperatures and in humid climates.

Do not use in conjunction with other sprays or fungicides.

If the crop to be treated has been subjected to freezing temperatures, do not treat until the crop has recovered before applying sulfur.

#### DIRECTIONS FOR USE

Always recommend use of 100% water for dilution. Use 100% water for dilution. Do not use less than 100% water for dilution. Do not use less than 100% water for dilution. Do not use less than 100% water for dilution. Do not use less than 100% water for dilution.

#### RECOMMENDATIONS

**APPLES:** Scab, powdery mildew. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Spray through second cover. Do not use sulfur on pink and red varieties. Use on sulfur-tolerant varieties only.

**CHERRY:** Blossom blight, brown rot, leaf spot, powdery mildew. Apply 4.5 lbs. per 100 gals. starting at shuck-fall.

**CITRUS:** Scab, blotch, and other diseases. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Apply 4.5 lbs. per 100 gals. (100 lbs. per acre).

**GRAPES:** Powdery mildew, leaf spot, rot. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Care should be taken to avoid surfaces from which the fungus is to be eradicated.

**PEAR:** Scab, powdery mildew, leaf spot, rot. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Care should be taken to avoid surfaces from which the fungus is to be eradicated.

**PEACH, NECTARINE:** Scab, leaf spot, rot. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Do not use sulfur on sensitive varieties. Do not use sulfur on sensitive varieties. Do not use sulfur on sensitive varieties.

**PLUM, PRUNE:** Scab, leaf spot, rot. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Do not use sulfur on sensitive varieties. Do not use sulfur on sensitive varieties.

**STRAWBERRIES:** Powdery mildew, leaf spot, rot. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Do not use sulfur on sensitive varieties. Do not use sulfur on sensitive varieties.

#### ORNAMENTALS

**ROSES, PERENNIALS, SHRUBS:** Powdery mildew, leaf spot, rot. Apply 4.5 lbs. per 100 gals. from petiole to pedicel. Do not use sulfur on sensitive varieties. Do not use sulfur on sensitive varieties.

**CAUTION:** Do not use sulfur on sensitive varieties. Do not use sulfur on sensitive varieties.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

50 LB. NET

**STAUFFER CHEMICAL COMPANY** NEW YORK, NY 10017