




# DUSTING SULFUR

## 98%



## AN AGRICULTURAL INSECTICIDE and FUNGICIDE

Active Ingredient:	
Sulfur .....	98%
Inert Ingredients .....	2%
	100%

### GUARANTEE OF FINENESS

All material contained in this bag is guaranteed to pass 95% through a 325 mesh screen. All screen tests are made in accordance with the California State Method, using screens certified by the United States Bureau of Standards.

### CAUTION

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on apricots, Anjou and comice pears, cranberries, cucurbits (cucumbers, squash and some varieties of melons), filberts, spinach, tung trees, walnuts and other sensitive plants.

Except on cotton, do not apply when shade temperatures exceed or are likely to exceed 90°F.

Sulfur may burn foliage and fruit during periods of high temperatures and under certain climatic conditions.

Do not use sulfur within 30 days of an oil spray (60 days on citrus).

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

**BURN EMPTY BAG—KEEP OUT OF SMOKE**

**50 LB. NET**

**STAUFFER CHEMICAL COMPANY**  
NEW YORK, NY 10017

### DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground power equipment or aircraft. Except as otherwise shown, apply at the first sign of infection and repeat at 7-10 day intervals as needed to maintain control.

### RECOMMENDATIONS

#### FRUIT CROPS

**Apples:** Scab, powdery mildew — Apply 30-35 lbs. per acre in the green tip to pink bud stage. Use on sulfur-tolerant varieties only.

**Cherry:** Blossom blight brown rot, powdery mildew — Apply 50 lbs. per acre in the popcorn, full bloom and petal fall stages; leaf spot, powdery mildew — Apply 50 lbs. per acre at petal fall, shuck-fall and 2 weeks after shuck-fall.

**Citrus:** Six-spotted and citrus rust mites — Apply 80 lbs. per acre; citrus flat and Yuma spider mites — Apply 100 lbs. per acre; citrus thrips — Apply 100 lbs. per acre in three applications; mid-March, petal-fall and 2-4 weeks later. Avoid applications during or preceding high temperatures. Do not apply within 2 months of an oil spray.

**Grapes:** Rust, powdery mildew — Apply 5-10 lbs. per acre in 4 applications: when shoots are 6" long; shoots 12-15" long; 14 days afterwards; 10-14 days thereafter to harvest.

**Plum, Prune:** Brown rot, leaf spot, prune rust — Apply 35-50 lbs. per acre starting 5 weeks before harvest and continuing at weekly intervals.

**Strawberries:** Powdery mildew — Apply 10-20 lbs. per acre.

#### VEGETABLE & FIELD CROPS

**Alfalfa:** Spider mites — Apply 15-25 lbs. per acre. Caution: sulfur may cause petal burn to open flowers.

**Beans:** Powdery mildew, rust, spider mites — Apply 15-30 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. Caution: May cause injury to some varieties of beans.

**Carrots:** Petroselin mite — Apply 30-35 lbs. per acre.

**Clover:** Powdery mildew, spider mites — Apply 15-25 lbs. per acre.

**Cotton:** Atlantic mite, cotton rust — Apply 25-30 lbs. per acre.

**Grass Seed Crops:** Timothy mite — Apply 30 lbs. per acre.

**Melon:** Honeydew and sulfur-resistant melons only) — Powdery mildew, spider mites — apply 20-25 lbs. per acre.

**Onions:** Petroselin mite — Apply 20-30 lbs. per acre during warm weather.

**Peas:** Powdery mildew — Apply 15-20 lbs. per acre.

**Sugar Beets:** Spider mite — Apply 30-35 lbs. per acre.

**Tomatoes:** Tomato russet mite — Apply 15-40 lbs. per acre.

**Turnips and Rutabagas:** Powdery mildew — Apply 15-20 lbs. per acre.

#### ORNAMENTALS:

Make full coverage applications on both sides of leaves. Keep new growth protected.

**Chrysanthemum:** Powdery mildew, rust. Do not apply when flowering.

**Hollyhock:** Rust.

**Rose, Pernetzia, Shrub:** Black spot, powdery mildew, rust. Caution: some burning may occur when used during warm weather.

USDA Reg. No. 476-2039

S-SF-690520-WC



# DUSTING SULFUR

ACCEPTED  
 May 29, 1969  
 UNDER U.S. GOVERNMENT  
 STANDARDS  
 476-2039

## 98%

### AN AGRICULTURAL INSECTICIDE and FUNGICIDE

Active Ingredient: Sulfur ..... 98%  
 Inert Ingredients ..... 2%  
 100%

#### GUARANTEE OF FINENESS

All material contained in this bag is guaranteed to pass 95% through a 325 mesh screen. All screen tests are made in accordance with the California State Method, using screens certified by the United States Bureau of Standards.

#### CAUTION

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on apricots, Anjou and comice pears, cranberries, cucurbits (cucumbers, squash and some varieties of melons), filberts, spinach, tung trees, walnuts and other sensitive plants.

Except on cotton, do not apply when shade temperatures exceed or are likely to exceed 90°F.

Sulfur may burn foliage and fruit during periods of high temperatures and under certain climatic conditions.

Do not use sulfur within 30 days of an oil spray (60 days on citrus).

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

**BURN EMPTY BAG—KEEP OUT OF SMOKE**

**50 LB. NET**

**STAUFFER CHEMICAL COMPANY**  
 NEW YORK, NY 10017

#### DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground power equipment or aircraft. Except as otherwise shown, apply at the first sign of infection and repeat at 7-10 day intervals as needed to maintain control.

#### RECOMMENDATIONS

##### FRUIT CROPS

**Apples:** Scab, powdery mildew — Apply 30-35 lbs. per acre in the green tip to pink bud stage. Use on sulfur-tolerant varieties only.

**Cherry:** Blossom blight, brown rot, powdery mildew — Apply 50 lbs. per acre in the popcorn, full bloom and petal fall stages; leaf spot, powdery mildew — Apply 50 lbs. per acre at petal fall, shuck-fall and 2 weeks after shuck-fall.

**Citrus:** Six-spotted and citrus rust mites — Apply 80 lbs. per acre; citrus flat and Yuma spider mites — Apply 100 lbs. per acre; citrus thrips—Apply 100 lbs. per acre in three applications; mid-March, petal-fall and 2-4 weeks later. Avoid applications during or preceding high temperatures. Do not apply within 2 months of an oil spray.

**Grapes:** Rust, powdery mildew — Apply 5-10 lbs. per acre in 4 applications: when shoots are 6" long; shoots 12-15" long; 14 days afterwards; 10-14 days thereafter to harvest.

**Plum, Prune:** Brown rot, leaf spot, prune rust—Apply 35-50 lbs. per acre starting 5 weeks before harvest and continuing at weekly intervals.

**Strawberries:** Powdery mildew — Apply 10-20 lbs. per acre.

##### VEGETABLE & FIELD CROPS

**Alfalfa:** Spider mites — Apply 15-25 lbs. per acre. Caution: sulfur may cause petal burn to open flowers.

**Beans:** Powdery mildew, rust, spider mites — Apply 15-30 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. Caution: May cause injury to some varieties of beans.

**Carrots:** Petrobia mite — Apply 30-35 lbs. per acre.

**Clover:** Powdery mildew, spider mites — Apply 15-25 lbs. per acre.

**Cotton:** Atlantic mite, cotton rust — Apply 25-30 lbs. per acre.

**Grass Seed Crops:** Timothy mite — Apply 30 lbs. per acre.

**Melon:** (Honeydew and sulfur-resistant melons only) — Powdery mildew, spider mites — apply 20-25 lbs. per acre.

**Onions:** Petrobia mite — Apply 20-30 lbs. per acre during warm weather.

**Peas:** Powdery mildew — Apply 15-20 lbs. per acre.

**Sugar Beets:** Spider mite — Apply 30-35 lbs. per acre.

**Tomatoes:** Tomato rust et mite — Apply 15-40 lbs. per acre.

**Turnips and Rutabagas:** Powdery mildew — Apply 15-20 lbs. per acre.

##### ORNAMENTALS:

Make full coverage applications on both sides of leaves. Keep new growth protected.

**Chrysanthemum:** Powdery mildew rust. Do not apply when flowering.

**Hollyhock:** Rust.

**Rose, Perennials, Shrubs:** Black spot, powdery mildew, rust. Caution: some burning may occur when used during warm weather.

USDA Reg. No. 476-2039

S-SF-690520-WC