


50 LB.



NET

COPPER-SULFUR 4-90 DUST

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Copper expressed as Metallic (from 5.33% Copper Oxide)	4.0%
Sulfur	90.0%
INERT INGREDIENTS	6.0%
	100.0%

CAUTION	Keep Out of Reach of Children
May cause irritation of nose, throat and skin. Do not breathe dust.	

OBSERVE THE FOLLOWING USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply to apricots, Anjou and Comice pears, cranberries, cucurbits (cucumber, cantaloupes, melons, squash), walnuts, or other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Do not use sulfur on any crop unless local use has proved safe.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing, number and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

Apply with specified amounts of this product uniformly to the foliage when air is calm using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5 to 10-day intervals, depending on severity of disease, but observe use limits given for specific crops.

When using ground equipment or aircraft apply at the rate of 20 to 40 lb. per acre, depending upon degree of infection and stage of plant development.

RECOMMENDATIONS

PEANUTS: Peanut Leaf Spot — Apply dust about 60 days after planting, or as soon as spots appear on leaf.

First Dust — Apply 15 to 20 lb. per acre.
Later Dusts — Apply 20 to 30 lb. per acre.

A minimum of three applications should be made at 2-week intervals.

Consult State Agricultural Experiment Station or State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.

ROSES: Black Spot, Powdery Mildew, Rust — Rose bushes should be dusted before the appearance of diseases on foliage. Apply at rate of 15 to 25 lb. per acre.

Repeat applications at weekly intervals until July (or until hot, dry weather begins); thereafter dust within 24 hours after each rain, but not more than once a week, until growth of plant is checked by cool weather.

BURN EMPTY BAG — KEEP OUT OF SMOKE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-2048 Made in U.S.A. T TA 6909I6 SB
Stauffer Chemical Company, New York

STAUFFER CHEMICAL COMPANY NEW YORK, NY 10017

50 LB.



12-9-69
476-2048

NET

COPPER-SULFUR 4-90 DUST

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Copper expressed as Metallic (from 5.33% Copper Oxide)	4.0%
Sulfur	90.0%
INERT INGREDIENTS	6.0%
	100.0%

CAUTION	Keep Out of Reach of Children
May cause irritation of nose, throat and skin. Do not breathe dust.	

OBSERVE THE FOLLOWING USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply to apricots, Anjou and Comice pears, cranberries, cucurbits (cucumber, cantaloupes, melons, squash), walnuts, or other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Do not use sulfur on any crop unless local use has proved safe.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing, number and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

Apply with specified amounts of this product uniformly to the foliage when air is calm using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5 to 10 day intervals, depending on severity of disease, but observe use limits given for specific crops.

When using ground equipment or aircraft apply at the rate of 20 to 40 lb. per acre, depending upon degree of infection and stage of plant development

RECOMMENDATIONS

PEANUTS: Peanut Leaf Spot - Apply dust about 60 days after planting, or as soon as spots appear on leaf.

First Dust - Apply 15 to 20 lb. per acre.
Later Dusts - Apply 20 to 30 lb. per acre.

A minimum of 3 to 4 applications should be made at 2-week intervals.

Consult State Agricultural Experiment Station or State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.

ROSES: Black Spot, Powdery Mildew, Rust - Rose bushes should be dusted before the appearance of diseases on foliage. Apply at rate of 15 to 25 lb. per acre.

Repeat applications at weekly intervals until July (or until hot, dry weather begins) thereafter dust within 24 hours after each rain, but not more than once a week, until growth of plant is checked by cool weather.

BURN EMPTY BAG - KEEP OUT OF SMOKE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-2048 Made in U.S.A. T TA 690916 SE
Stauffer Chemical Company, New York

STAUFFER CHEMICAL COMPANY NEW YORK, NY 10017