

TRITHION[®]-SEVIN-SULFUR 3-5-30 DUST

3-1-72
476-2066

AGRICULTURAL ACARICIDE-INSECTICIDE-FUNGICIDE

ACTIVE INGREDIENTS:

Carbophenothion: S-(p-chlorophenylthio)methyl O,O-diethyl phosphorodithioate**	3%
Carbaryl: (1-naphthyl N-methylcarbamate)	5%
Sulfur	30%
INERT INGREDIENTS:	62%
	<hr/> 100%

*Trade-mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate
**U.S. Pat. No. 2,793,224

CAUTION

Keep Out of Reach of Children

HAZARDOUS IF SWALLOWED OR INHALED.

Avoid breathing dust, contact with skin and eyes.

Wear dust respirator and goggles.

Wear regular long-sleeved work clothing. Change to clean clothing daily.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

Take shower or bath after work.

Wash all contaminated clothing with soap and hot water before reuse.

Do not contaminate food, feed, feed troughs and watering receptacles.

NOTE TO PHYSICIAN

Trithion[®] can cause cholinesterase inhibition. ATROPINE IS ANTI-DOTAL.

BURY WASTE AND EMPTY CONTAINER IN A SAFE PLACE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

E.P. Reg. No. 476-2066 AA

Made in U.S.A. by
STAUFFER CHEMICAL COMPANY
Westport, Ct. 06980

50 LB. NET

OBSERVE THE FOLLOWING USE PRECAUTIONS

Read and follow carefully all directions with all label directions.

Sulfur will cause severe frost and water injury to sulfur sensitive crops. Do not use sulfur on any crop unless weather forecast indicates that it may burn foliage and fruit during periods of high temperature and low humidity. Other climatic conditions. Do not apply sulfur to sensitive crops when high temperature and humidity are likely to exceed 90° F.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

In order that pesticides do not harm and damage crops will not exceed tolerance established by the Food and Drug Administration, apply at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or their animals, or onto adjoining food, fiber or pasture crops. **The grower is responsible** for the damage to his crops as well as to damages caused by drift from his property to that of others.

FISH AND WILDLIFE CAUTIONS: Toxic to fish and wildlife. Do not contaminate any body of water. Do not apply to any stream or pond on this water. Do not contaminate water by disposal in water. Birds, fish and other animals may be killed.

RECOMMENDATIONS AND DIRECTIONS FOR USE

This material is to be applied to crops and plants by hand using a backpack sprayer or other suitable equipment.

BEANS, DRY (SHELLED): Bean aphid, bean web beetle, common bean beetle, Mexican bean beetle, Mexican bean beetle, potato leafhopper, two-spotted mite, serpentine leaf miner, velvetbean caterpillar and Western bean cutworm. Apply 10 and 15 lbs. per acre. Do not apply more than two applications per season. Do not use on or near any water within 1 day after application.

BEAN, SNAP: Bean aphid, serpentine leaf miner, potato leafhopper, Mexican bean beetle, two-spotted mite, velvetbean caterpillar, and cutworm. Apply within 7 days of harvest. Do not apply more than two applications per season. Do not use on or near any water within 1 day after application.

GRAPE: Grape leafhopper, grape leafhopper, sawfly, mildew spider mite, two-spotted mite, and grape leaf miner. Apply 10 and 15 lbs. per acre. Do not apply more than one application per season. Make the application at dusk, with a uniform spray. If the pest is not controlled, a second application may be made immediately after the first application. Do not use on or near any water within 1 day after application. Do not use on or near any water within 1 day after application. Do not use on or near any water within 1 day after application.

PEAS: Pea aphid, potato leafhopper, and serpentine leaf miner. Apply 10 and 15 lbs. per acre. Do not use on or near any water within 1 day after application.

PEAS: Pea aphid, potato leafhopper, and serpentine leaf miner. Apply 10 and 15 lbs. per acre. Do not use on or near any water within 1 day after application.