


ELECTRIC[®] SULFUR-ZINC DUST

(92.5-4.82)

AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENT:

Sulfur 92.5%

INERT INGREDIENTS* 7.5%

*Also contains 4.82% Zinc expressed as metallic
(from 6% Zinc Oxide). 100.0%

CAUTION

Keep Out of Reach of Children

® Registered Trademark of Stauffer Chemical Co.

BURN EMPTY BAG — KEEP OUT OF SMOKE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-2073
ARC 710222

Made in U.S.A.
Stauffer Chemical Company
New York, New York 10017

50 LB. NET

OBSERVE THE FOLLOWING USE PRECAUTIONS

USE ONLY ON CROP LISTED.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90°F.

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply or allow to drift to adjoining crops. The grower is responsible for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service, as the timing, number and rate of applications needed will vary with local conditions.


RECOMMENDATIONS AND DIRECTIONS FOR USE

GRAPES: For the control of powdery mildew and for correction of zinc deficiency (little leaf), apply 10 to 30 lbs. per acre in each of four applications. Make first application when shoots are 6 inches long, second when shoots are 12 to 15 inches long, third 14 days later and the last 10 to 14 days after the third application.

ELECTRIC[®] SULFUR—ZINC DUST

(92.5-4.82)

AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENT:

Sulfur 92.5%

INERT INGREDIENTS*

7.5%

*Also contains 4.82% Zinc expressed as metallic
(from 6% Zinc Oxide).

100.0%

CAUTION

Keep Out of Reach of Children

[®] Registered Trademark of Stauffer Chemical Co.

BURN EMPTY BAG — KEEP OUT OF SMOKE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-073
ARC 710222

Made in U.S.A.
Stauffer Chemical Company
New York, New York 10017

50 LB. NET

OBSERVE THE FOLLOWING USE PRECAUTIONS

USE ONLY ON CROP LISTED.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90 F.

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply or allow to drift to adjoining crops. The grower is responsible for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service, as the timing, number and rate of applications needed will vary with local conditions.

RECOMMENDATIONS AND DIRECTIONS FOR USE

GRAPES: For the control of powdery mildew and for correction of zinc deficiency (little leaf), apply 10 to 30 lbs. per acre in each of four applications. Make first application when shoots are 6 inches long, second when shoots are 12 to 15 inches long, third 14 days later and the last 10 to 14 days after the third application.