

CAPTAN-THIODAN 7.5-3 DUST

AGRICULTURAL INSECTICIDE AND FUNGICIDE

ACTIVE INGREDIENTS

Captan - N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

7.5%

Endosulfan*

3.0%

INERT INGREDIENTS:

89.5%

100.0%

*Endosulfan is a registered trademark of Shell Chemical Co.

Thiodan is a registered trademark of Canadian Home Products Co.

Thiodan is a registered trademark of Canadian Home Products Co. Patent No. 1,199,887

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN

Avoid breathing dust.

Avoid contact with skin, eyes or clothing.

In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention.

Wash with soap and water after handling and before eating, or smoking; wear clean clothing.

Avoid contamination of feed and food.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not apply to plants which are stressed or which have been damaged by frost, drought, or other adverse conditions.

Do not apply to plants which are in bloom or which are about to bloom.

Do not apply to plants which are in fruit or which are about to fruit.

The grower is responsible for the proper use of this product.

Do not apply to plants which are in fruit or which are about to fruit.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops.

For the control of disease persisting after the prescribed time limits have expired, an effective fungicide should be used alone in a manner which will not leave residue in excess of established tolerance.

CAUTION: Workers entering treated areas the day of application should wear protective clothing.

RECOMMENDATIONS

CUCUMBER, PUMPKIN, SUMMER & WINTER SQUASH: Angular leaf spot, anthracnose, downy mildew, aphids, cucumber beetle, striped cabbage flea beetle, squash vine borer. Apply 33 lb. per acre. For squash vine borer control, apply weekly to flower buds, stems and vines beginning when mites first appear. May be applied up to day of harvest.

POTATOES: Early & late blight, aphids, European corn borer, Colorado potato beetle, false chinch bug, flea beetle, green stink bug, leafhopper, leaf-footed plant bug, potato psyllid, potato tuberworm, three-lined potato beetle, Southern armyworm. Apply 3-6 lb. per acre. Repeat as needed.

STRAWBERRIES: Strawberry root, yellow spittlebug, strawberry aphid. Apply 33 lb. per acre. Do not apply within 4 days of picking. Do not apply at intervals of less than 15 days or more than twice during any 30-day period if fruit is present.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the accuracy of merchantability, and is liable for any particular purpose concerning this product, except as stated on the label.

NO. 10-11-11

BURN EMPTY BAG—KEEP OUT OF SMOKE