

CAPTAN-HEXACHLOROBENZENE 20-20 SEED PROTECTANT

FOR SEED DISEASE CONTROL

SPECIAL LUBRICANT ADDED - DISCOLORED

ACCEPTED
MAR 19 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 476-2070

ACTIVE INGREDIENTS

Captan: N-[(Trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide*	20%
Hexachlorobenzene	20%
INERT INGREDIENTS	60%
	100%

*Not licensed for use in countries where valid patients still exist.

CAUTION

Keep Out of Reach
of Children

HARMFUL IF TAKEN INTERNALLY.

Avoid inhaling dust, wear respirator.
Avoid contact with skin, eyes and clothing.
Wash with soap and water after handling.
Avoid contamination of food and feed.

GENERAL USE PRECAUTIONS

- Read all directions before using and use only as specified on this label.
- Toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Dispose of excess treated seeds by burial.
- **IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED
DO NOT USE FOR FOOD, FEED OR OIL PURPOSES

The federal government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

PROTECT FROM EXCESSIVE HEAT—STORE IN A COOL DRY PLACE.

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES.

Made in U.S.A.
Stauffer Chemical Company
New York, N.Y. 10017

RECOMMENDATIONS

Captan-Hexachlorobenzene 20-20 Seed Protectant is a special formulation for the treatment of wheat seeds prior to planting for the protection of the seeds against bunt (stinking smut), seed rots, seedling blights and root rot. It can be used by the planter box method or by dust treatment.

DIRECTIONS FOR USE

FOR PROTECTION OF WHEAT SEED FROM BUNT (STINKING SMUT),
SEED ROTS, SEEDLING BLIGHTS AND ROOT ROT

PLANTER BOX TREATMENT: 2 ounces per bushel seed (3 1/3 ounces per 100 pounds seed). Mix thoroughly with seed in planter box before planting.

DUST TREATMENT: Apply 2 ounces per bushel of seed (3 1/3 ounces per 100 pounds seed). Mix thoroughly with seed in dusting machine, barrel or other container. May also be applied through drill box or planter box.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-2070

S-ARC-710212-MW

50 LB. NET