



CAPTAN-THIRAM-METHOXYCHLOR 42-42-1.5 SEED PROTECTANT

DYE ADDED

SLURRY APPLICATION FOR SEED DISEASE AND INSECT CONTROL

OBSERVE THE FOLLOWING USE PRECAUTIONS

READ ALL LABEL DIRECTIONS BEFORE USING
Use only for claims listed and only as recommended on this label.

IMPORTANT: Treated seed must not be used for human consumption or as animal feed. Bag containing treated seed should be labeled:

TREATED SEED.

DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.

The Federal Government and some states have issued regulations for the application of special taxes which apply to be affixed to bags containing treated seed.

FISH CAUTIONS. Do not permit treated seed to come in contact of any body of water. Do not use materials which are cleaned of equipment in contact with water.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

ACTIVE INGREDIENTS:	
Captan*	42.0%
Thiram (tetramethylthiuram disulfide)	42.0%
Methoxychlor, Technical**	1.5%
INERT INGREDIENTS:	14.5%
	<hr/> 100.0%

* Captan is a trademark of ICI, Inc. (Ciba-Geigy) Ltd. dicarboximide. Not to be used in countries where seed patents are in effect.
** Equivalent to 1.5% of the active methoxyphenyl-1,1,1-trichloro-2,2,2-trifluoro-2-methyl-2-propanol and related compounds.

CAUTION Keep Out of Reach of Children

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.

Do not inhale dust or vapors.
Do not get on face, hands, or clothing.
Do not get on skin or clothing.
Do not get on eyes.
Do not get on mouth.

50 LB. NET

DIRECTIONS FOR USE

Slurry Treatment: Pre-mix the recommended amount of this material with a little water in a bucket and make a thick paste. Stir this paste into the required amount of water for the slurry equipment to be used. **Note:** Seed treated by the slurry method should not be sown until such time as the seed has had an opportunity to dry. Store in well ventilated area. If seed is stored in a wet, humid, or damp environment, it may reduce germination.

RECOMMENDATIONS

For control of many seed rots, seedling rot, root rots, fungi, and for protection of stored seed against weevils, confused flower beetle, flat grain beetle, for grain beetle, granary weevil, hairy fungus beetle, lesser grain borer, long-headed flour beetle, red flour beetle, rice weevil, and saw-toothed grain beetle.

AMOUNT OF CAPTAN-THIRAM-METHOXYCHLOR 42-42-1.5 TO USE ON SEEDS

SEED	SLURRY
Wheat	1.0-2.0%
Barley	1.0-2.0%
Oats	1.0-2.0%

Further details of treatment of seeds may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

PROTECT FROM EXCESSIVE HEAT • STORE IN A COOL DRY PLACE
CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES

STAUFFER CHEMICAL COMPANY

 NEW YORK, N.Y. 10017