



# CAPTAN-THIRAM

## 43-43 SEED PROTECTANT

DYE ADDED

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT AND ECONOMIC POISON CONTROL ACT UNDER NO. 100-2075

### SLURRY APPLICATION AND DRY TREATMENT FOR SEED DISEASE CONTROL

#### ACTIVE INGREDIENTS:

Captan: N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide 43.0%  
Thiram (tetramethylthiuram disulfide) 43.0%

#### INERT INGREDIENTS:

14.0%

100.0%

### CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.

AVOID INHALING DUST — WEAR A RESPIRATOR.

AVOID CONTACT WITH SKIN. WASH WITH SOAP AND WATER AFTER HANDLING.

CONSUMPTION OF ALCOHOLIC BEVERAGES INCREASES THE POSSIBILITY OF HARM.

DO NOT CONTAMINATE FOOD OR FEED.

PROTECT FROM EXCESSIVE HEAT — STORE IN A COOL DRY PLACE.

**BURY WASTE AND EMPTY CONTAINER IN AN ISOLATED AREA AWAY FROM WATER SUPPLIES.**

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

50 LBS. NET

EPA Reg. No. 476-2093  
EPA Est. 476-NB-1

A-1

Made in U.S.A. By

# STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

#### OBSERVE THE FOLLOWING USE PRECAUTIONS

READ THE LABEL DIRECTIONS BEFORE USING. Use only for claims listed and only as recommended on this label.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Dispose of excess treated seed by burial. Bags containing treated seed should be labeled:

**TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.**

The federal government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

FISH CAUTIONS: This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE AND RECOMMENDATIONS

##### SLURRY TREATMENT

Pre-mix the recommended amount of this material with a little water in a bucket and make thin paste. Stir this paste into the required amount of water for the slurry equipment to be used.

NOTE: Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas.

For control of seed and seedling diseases, damping-off, seed decay, seedling blights, seed rots, molds, storage rots, covered smuts of barley and wheat, loose and covered smuts of sorghum, seed and soil-borne fungi.

##### AMOUNT OF CAPTAN-THIRAM 43-43 SEED PROTECTANT TO USE ON SEEDS

| SEED                                    | Amount (oz./cwt.) |
|-----------------------------------------|-------------------|
| Barley                                  | 4.50              |
| Beans (Snap and Dry), Soybeans, Cowpeas | 2.50              |
| Corn (Field)                            | 2.60              |
| Corn (Sweet)                            | 8.75              |
| Cotton                                  | 5.00              |
| Peas, Beans (Large Lima), Sesame        | 3.75              |
| Safflower                               | 4.50              |
| Sorghum, Millet                         | 5.00              |
| Tomatoes                                | 2.00              |
| Wheat                                   | 4.00              |

##### DUST TREATMENT

Mix thoroughly in a dusting machine, barrel or other effective dry mixer.

|                       |       |
|-----------------------|-------|
| Beets (Table)         | 10.00 |
| Cantaloupe, Cucumbers | 5.50  |
| Muskmelon, Watermelon | 3.50  |
| Peanuts (Shelled)     | 4.50  |
| Peanuts (Unshelled)   | 4.00  |
| Pumpkin, Squash       | 4.00  |
| Spinach               | 9.50  |

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.