


APTA 43-43 SEED PROTECTAN.



SLURRY APPLICATION AND DRY TREATMENT FOR SEED DISEASE CONTROL

ACTIVE INGREDIENTS: Captan: N-I (trichloromethyi)thiol-4-cyclohexene-1,2 dicarboximide 2.34% .. 43.00% Related derivatives
Thiram (tetramethylthiuram disulfide) INERT INGREDIENTS: 14.00% 100.00% Keep Out of Reach of Children

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.
AVOID INHALING DUST — WEAR A RESPIRATOR.
AVOID CONTACT WITH SKIN. WASH WITH SOAP AND WATER AFTER HANDLING.
CONSUMPTION OF ALCOHOLIC BEVERAGES INCREASES THE POSSIBILITY OF HARM.
DO NOT CONTAMINATE FOOD OR FEED.

PROTECT FROM EXCESSIVE HEAT - STORE IN A COOL DRY PLACE.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE disposal. Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for posticides or buried in a safe place have yet meaning supplies.

away from water supplies containers disposals. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Generally in a safe place. Generally in a safe place. Generally in a safe place generally in a safe place or local disposal authorities for approved alternative procedures such as limited open burning.

NOTICE - READ CAREFULLY

CONDITIONS OF GALE S. suffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION. Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to it. The directions for use on the label subject to the inherent risks referred to below. Stauffer makes no ower express warranties, THERE IS NO IMPLIED WARRANTY OF IMERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY in no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the gurchase price of tra groduct in respect of which such damages are claimed.

This product is said only for uses a 'ded on the label. No express or implied license is granted to use or sell this product under any patent' any country except as specified: Country: U.S.A.

POUNDS NET

EPA Reg. No. 476-2053

EFA Est. No. 476- -

A-2

OBSERVE THE FOLLOWING USE PRECAUTIONS

READ THE LABEL DIRECTIONS BEFORE USING. Use only for claims listed and only as recommended on this label.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Dispose of excess treated seed by burial. Bags containing treated seed should be labeled:

treated seed. Do not use for food, feed or oil purposes. The federal government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

FISH CAUTIONS: This product is toxic to lish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE AND RECOMMENDATIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SLURRY TREATMENT

Pre-mix the recommended amount of this material with a little water in a bucket and make thin paste. Stir this paste into the required amount of water for the slurry equipment to be used. NOTE: Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas.

For control of seed and seedling diseases, damping-off, seed decay, seedling blights, seed rots, molds, storage rots, covered smuts of barley and wheat, loose and covered smuts of sorghum, seed and soil-borne fungi.

AMOUNT OF CAPTAN-THIRAM 43-43 SEED PROTECTANT TO USE ON SEEDS SEED

Barley	4.50 oz./	cwt.
Beans (Snap and Dry), Soybeans, Cowpeas	2.50 oz./	cwt.
Com (Field)	7.60 nz. i	rewr
Corn (Sweet)	8.75 oz./	cwt.
Cetton	5.00 oz./	CWI.
Peas, Beans (Large Lima), Sesame	3.75 OZ./	cwt.
Safflower	4.50 oz./	cwt.
Sorghum, Millet	5.00 oz./	cwt.
Tematoes	2.00 oz./	cwt.
Wheat	4.00 02./	CWL
DUST TREATMENT		

Mix thoroughly in a dusting machine, barrel or other effective dry mixer.

Beets (Table)
Cantaloupe, Cucumbers
Muskmelon, Watermelon
Peanuts (Shelled)
Peanuts (Unshelled)
Pumpkin, "quash
Spinach

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

Made In U.S.A. By STAUFFER CHEMICAL COMPANY