

6-11-71
476-2096

TRITHION-SEVIN[®] 3-5 DUST

AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:

Carbophenothion S-[(p-chlorophenylthio)methyl] O,O-diethyl phosphorodithioate**	3%
Carbaryl (1-naphthyl N-methylcarbamate)	5%

INERT INGREDIENTS:

92%
100%

**U.S. Pat. No. 2,793,224

*Trade-mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

CAUTION

**Keep Out of Reach
of Children**

HAZARDOUS IF SWALLOWED OR INHALED.

Avoid breathing dust, contact with skin and eyes.

Wear dust respirator and goggles.

Wear regular long-sleeved work clothing. Change to clean clothing daily.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

Take shower or bath after work.

Wash all contaminated clothing with soap and hot water before re-use.

Do not contaminate food, feed, feed and troughs and watering receptacles.

NOTE TO PHYSICIAN

Trithion-Sevin upon repeated or prolonged use may cause cholinergic inhibition. ATROPINE IS ANTIDOTAL.

BURY EMPTY CONTAINER AND WASTE IN A SAFE PLACE

OBSERVE THE FOLLOWING USE PRECAUTIONS

Use only for claim listed and in accordance with all label directions.

Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on this label. Do not contaminate water by disposal of wastes. Birds feeding on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative agricultural extension service.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Extension Service or State Agricultural Experiment Station in your region for specific information as to timing, rate and number of applications. Always vary with local conditions.

RECOMMENDATIONS AND DIRECTIONS FOR USE

This formulation is to be applied only by experienced personnel using approved safety equipment.

GRAPE: Grape leaf folder, grape vine leafhopper, grape vine trunk borer, grape vine and Western grape leaf skeletonizer.

Apply 25 to 30 pounds per acre by air-blast or ground equipment when air is calm. Make thorough coverage of plants with a uniform deposit of dust. Repeat as needed. For grape vine leafhopper control, apply immediately after a majority of leafhoppers have hatched. Thorough under leaf coverage is important.

Do not apply within 10 days of harvest. Do not apply to sensitive varieties such as Chardonnay, Gamay, Golden Muscat, Muscat and Italia.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

50 LB. NET