



CAPTAN-METHOXYCHLOR 75-5 SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE AND INSECT CONTROL

DYE ADDED

CAPTAN-METHOXYCHLOR 75-5 SEED PROTECTANT is a special wettable powder formulation for the slurry treatment of seeds, prior to planting, for protection against certain fungus diseases and insects.

GENERAL USE PRECAUTIONS

Read All Directions Before Using and Use Only as Specified on This Label.

Toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of excess treated seeds by burying with wastes.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED

DO NOT USE FOR FEED, FOOD OR OIL PURPOSES
The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

DIRECTIONS FOR USE

SLURRY SEED TREATMENT: Pre-mix the recommended amount of this material with a little water in a bucket and make a thin paste. Stir this paste into the required amount of water for the slurry equipment to be used.

NOTE: Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas. If seed is bagged when wet, heat is developed and impairs seed germination.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

ACTIVE INGREDIENTS:

Captan*	75.0%
Methoxychlor, Technical**	5.0%

INERT INGREDIENTS:	20.0%
	100.0%

*N-[[trichloromethyl) thio]-4-cyclohexene-1, 2-dicarboximide.

Not licensed for use in countries where valid patents still exist.

**Equivalent to 4.4% of 2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 0.6% other isomers and related compounds.

U. S. Pat. No. 2,420,928.

CAUTION

Keep Out of
Reach of Children

Harmful if swallowed, inhaled, or absorbed through skin.

Avoid inhaling dust—wear respirator.

Avoid contact with skin.

Wash with soap and water after handling.

Do not contaminate feed or food.

50 LB. NET

RECOMMENDATIONS

For control of many seed rots, seedling rots, root rots fungi, kernel smut of sorghum, and for protection of stored seed against Cadelle, Confused Flour Beetle, Flat Grain Beetle, Granary Weevil, Lesser Grain Borer, Red Flour Beetle, Rice Weevil, and Saw-Toothed Beetle.

AMOUNT OF CAPTAN-METHOXYCHLOR 75-5 TO USE ON SEEDS

SEED	SLURRY
Alfalfa, Clover, Lespedeza, Trefoil.....	5½ oz./100 lb.
Barley, Oats, Wheat.....	2½ oz./bu.
Beans (dry, snap, lima), Cowpeas, Peas.....	1 oz./bu.
Beets (garden).....	8 oz./100 lb.
Cantaloupe, Cucumber.....	2 oz./100 lb.
Cole crops (broccoli, Brussels sprouts, cabbage, cauliflower).....	1 oz./100 lb.
Corn — Field.....	2½ oz./bu.
Sweet.....	2½ oz./100 lb.
Cotton — Acid delinted.....	2 oz./100 lb.
Fuzzy.....	3 oz./100 lb.
Crucifers.....	
(Mustard, Radishes, Rape, Turnips).....	1 oz./100 lb.
Flax.....	1½ oz./bu.
Grasses.....	5½ oz./100 lb.
Muskmelon, Pumpkin, Squash, Watermelon.....	1½ oz./100 lb.
Sorghum (hull-less).....	1½ oz./bu.
Soybeans.....	1 oz./bu.
Spinach.....	4 oz./100 lb.
Sugar beets — West.....	4 oz./100 lb.
East.....	8 oz./100 lb.
Swiss chard.....	8 oz./100 lb.

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

EPA REG. NO. 476-2102

Made in U.S.A.

ARC-710827-0

PROTECT FROM EXCESSIVE HEAT • STORE IN A COOL DRY PLACE

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES

STAUFFER CHEMICAL COMPANY NEW YORK, NY 10017

476-2102
9/15/1971
122

ACCEPTED

SEP 15 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

476-2102