



CAPTAN-METHOXYCHLOR 75-5 SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE AND INSECT CONTROL

DYE ADDED

CAPTAN-METHOXYCHLOR 75-5 SEED PROTECTANT is a special wettable powder formulation for the slurry treatment of seeds, prior to planting, for protection against certain fungus diseases and insects.

GENERAL USE PRECAUTIONS

Read All Directions Before Using and Use Only as Specified on This Label.

Toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of excess treated seeds by burying with wastes.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled.

TREATED SEED

DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

The Federal Government and some state have regulations for the application of special labels to be affixed to bags containing treated seed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SLURRY SEED TREATMENT: Premix the recommended amount of this material with a little water in a bucket and make a thin paste. Stir this paste into the required amount of water for the slurry application to be used.

NOTE: Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas. If seed is bagged when wet heat is developed and impairs seed germination.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide spray mixture or residue that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

ACTIVE INGREDIENTS:

CAPTAN: N-(1-trichloromethyl)thio-4-cyclohexene 70.92%

1,2 dicarboximide 4.08%

Related derivatives 5.00%

Methoxychlor 20.00%

INERT INGREDIENTS:

100.00%

Equivalent to 4.4 lb. of 2,2-bis(4-methoxyphenyl)-1,1,1-trichloroethane and 0.6 lb. other isomers and related compounds

CAUTION Keep Out of Reach of Children

Harmful if swallowed, inhaled, or absorbed through skin.

Avoid inhaling dust — wear respirator.

Avoid contact with skin.

Wash with soap and water after handling.

Do not contaminate feed or food.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE: Stauffer (and sealer) offers this product for sale to you and you and all users are deemed to have accepted the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION: Stauffer warrants that this product conforms to the form or description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties. THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and the claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

EPA Reg. No. 476-107

EPA Est. 476-NB-1

50 LB. NET

PROTECT FROM EXCESSIVE HEAT • STORE IN A COOL DRY PLACE

RECOMMENDATIONS

For control of many seed rots, seedling rots, root rots fungi, kernel smut of sorghum, and for protection of stored seed against Cadelle, Confused Flour Beetle, Flat Grain Beetle, Granary Weevil, Lesser Grain Borer, Red Flour Beetle, Rice Weevil, and Saw-Toothed Beetle.

AMOUNT OF CAPTAN-METHOXYCHLOR 75-5 TO USE ON SEEDS

SEED	SLURRY
Alfalfa, Clover, Lespedeza, Trefol	5 1/2 oz./100 lb.
Barley, Oats, Wheat	2 1/2 oz./bu.
Beans (dry, snap, lima), Cowpeas, Peas	1 oz./bu.
Beets (garden)	8 oz./100 lb.
Cantaloupe, Cucumber	2 oz./100 lb.
Cole crops (broccoli, Brussels sprouts, cabbage, cauliflower)	1 oz./100 lb.
Corn - Field	2 1/2 oz./bu.
Sweet	2 1/2 oz./100 lb.
Cotton - Acid delinted	2 oz./1 lb.
Fuzzy	3 oz./1 lb.
Crucifer -	
Mustard, Radishes, Rape, Turnip	1 oz./100 lb.
Flax	1 1/2 oz./bu.
Grasses	5 1/2 oz./100 lb.
Muskmelon, Pumpkin, Squash	
Watermelon	1 1/2 oz./100 lb.
Sorghum (dry, sweet)	1 1/2 oz./bu.
Soybeans	1 oz./bu.
Spinach	4 oz./100 lb.
Sugar beets - West	4 oz./100 lb.
East	8 oz./100 lb.
Swiss chard	8 oz./100 lb.

For further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

This product is sold only for uses stated on the label. No express or implied warranty is granted by use of this product under any patent in any country except as specified. Country: USA

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880