GENERAL USE PRECAUTIONS

Read all label instructions before using. Do not overdose. ORDRAM should be used only for recommended purposes and at recommended rates. ORDRAM may cause crop injury under extreme soil or climatic conditions or if directions are not followed.

Except as specified on this label, do not apply other herbicides in conjunction with <code>RDRAM</code> as injury may occur.

ORDRAM may be combined with fluid (solution, slurry, or suspension) fertilizers. However, physical compatibility with these fluid fertilizers must be determined before combining in the spray tank. See special literature giving directions for these combinations. Even though found to be compatible, constant agitation is necessary to keep the ORDRAM uniformly mixed with the fluid fertilizer.

DO NOT STORE NEAR SEEDS OR FERTILIZER.

KEEP CONTAINER CLOSED WHEN NOT IN USE.

DO NOT CONTAMINATE WATER USED FOR DOMESTIC PURPOSES.

GENERAL APPLICATION DIRECTIONS

All equipment used in the application of ORDRAM should be carefully calibrated before use and checked frequently during application to be sure the application is uniform. Avoid overlaps that will increase ORDRAM dosages above recommended limits because crop injury may occur. Apply ORDRAM 8-E in 10 to 50 gallons of water by ground equipment and 5 to 10 gallons of water by airplane except as specified below.

To avoid uneven distribution of ORDRAM in flooded fields the water flow through the field should be at a minimum at the time of application and for at least two hours after application.

Do not apply more than 6 pounds actual ORDRAM (6 pints ORDRAM 8-E) per acre per season where more than one method of application is used. Allow 30 days between applications.

DIRECTIONS FOR USE AND RECOMMENDATIONS

 PRE-PLANT, PRE-FLOOD: For use on waterseeded rice ONLY.

For control of Watergrass (Barnyardgrass) Echinochioa spp. ONLY.

SOIL PREPARATION: Prepare the soil for seeding according to good agricultural practices. All weed growth must be thoroughly worked into the soil before treatment.

FOR GROUND APPLICATION ONLY: Use standard low-pressure (20-50 psi) boom-spray equipment. DO NOT USE SINGLE NOZZLE BOOM-JET TYPE SPRAYERS. APPLICATION AND INCORPORATION: Apply to dry soil in good tilth. ORDRAM must be incorporated immediately (within minutes) to prevent loss of the herbicide. Whenever possible, application and incorporation should be done in the same operation. Incorporation should be done by at least one discing or spiketooth harrowing. Flood the field for seeding and hold the flood for 4 to 6 days following seeding.

RECOMMENDED RATE: Apply 3 pints ORDRAM 8-E per acre.

POST-EMERGENCE, PRE-FLOOD: For use on dry-seeded and water-seeded rice.

For control of Dayflower (Commelina communis) in the two leaf stage or less and small Sprangletop (Leptochloa spp.) emerged less than one inch, apply ORDRAM 8-E as a tank mix combination with a registered propanil brand in 10-15 gallons of water per acre by airplane or in 20-50 gallons of water per acre by ground equipment. Flood the rice within 5 days after treatment. Follow all directions and use precautions on the propanil label. Avoid drift or accidental application to all crops, shrubs or ornamentals not in the rice field being treated.

RECOMMENDED RATE: 2 to 3 pints ORDRAM 8-E (3 pints for Sprangletop) plus 3 pounds actual propanil per acre.



Ordram

An Emulsifiable Liquid Herbicide for Use on Rice THIS LABEL CONTAINS DIRECTIONS FOR USE ON RICE GROWN IN THE SOUTHWEST ONLY ACTIVE INGREDIENT:

S-ethyl hexahydro-1H-azepine-1-carbothioate
INERT INGREDIENTS

90.9% 9.1% 100.0%

Contains 8 Pounds Active Ingredient Per Gallon.

U. S. Pat. No. 3.198.786

WARNING

Keep Out of Reach of Children

May be fatal if swallowed. Avoid contact with skin, eyes and clothing. Wash with soap and water after use. Avoid breathing spray mist. Avoid contamination of feed and food. Repeated skin contact may cause cumulative effects.

EPA Reg. No. 476-2107

Made In U.S.A. By

GALLONS NET

STAUFFER CHEMICAL COMPANY WESTPORT, CT 06880

3. POS dry-s For flow

For flow Whe reco

head REC

tion Bari Floc unti

4. **PO**: dry-

to to only

The but 3 s (no**t**

Wat rice



Ordram

An Emulsifiable Liquid Herbicide for Use on Rice
THIS LABEL CONTAINS DIRECTIONS FOR
USE ON RICE GROWN IN THE SOUTHWEST ONLY
ACTIVE INGREDIENT:

S-ethyl hexahydro-1H-azepine-1-carbothioate INERT INGREDIENTS

90.9% 9.1%

100.0%

Contains 8 30 ands Active Ingredient Per Caden

use on water

) Echinophinic

or seeding ac-

ll weed growth.

before treat-

standard lowent. DO NOT

E SPRAYERS

Apply to dry

orporated 🚓

t loss of the

on and are to

eration, incor

ne discong or

r seeding and

ORDRAM 3-E

: For use on

nmunis) in the

p*(Leptochica* ORDRAM & E

tered propanii

by airplane or

nd equipment

tment Follow

the propanilication to all

the rice field.

RDRAM 81 (3)

ctual propanii

seeding.

U. S. Pat. No. 3 198 736

WARNING Keep Out of Reach of Children

May be fatal if swallowed. Avoid contact with skin, eyes and clothing. Wash with soap and water after use. Avoid breathing spray mist. Avoid contamination of feed and food. Repeated skin contact may cause cumulative effects.

EPA Reg. No. 476-2107

GALLONS NET

STAUFFER CHEMICAL COMPANY
WESTPORT, CT 06880

Made in U.S.A. By

3 **POST-EMERGENCE AT FLOODING:** For use on dry-seeded rice ONLY.

For control of Watergrass (Barnyardgrass) and Day flower under 5 inches tall.

When establishing the permanent flood, meter the recommended rate of ORDRAM 8-E per acre into the irrigation water at the header ditch or canal where water enters the treated field. Meter the ORDRAM during the entire irrigation period using a constant head metering device.

RECOMMENDED RATE: 3 pints ORDRAM 8 El per activi

The rice must be established at the time of application. Dayflower must be completely submerged and Barnyardgrass 23 submerged by the flooding operation. Flood should be maintained at the prescribed depth until Barnyardgrass and or Dayflower is controlled.

 POST-EMERGENCE, POST-FLOOD: for use on dry-sheded and water seeded rich.

For control of Watergrass (Barnyardgrass, ONLY).

APPLICATION: Apply the recommended rate directly to the flooded field. Apply by airplane ONLY. Apply only when wind velocity is low, 0 to 5 mpi.

RECOMMENDED RATE: 3 pints ORDRAM 8-E per a re-

The Watergrass must be a minimum of 2 inches high but not higher than 5 inch s, and it must be at least 23 submerged at time of application. This water level (not less than 2 inches) must be maintained until the Watergrass dies. Application should be made before rice is 2 inches above surface of water

NOTICE: Stauffer Chemical Compa owarranty, express or implied, including the war, es of merchantability, and/or fitness for an particular purpose, concerning this material, except those which are contained on Stauffer's label.


DISPOSAL:

RINSE SPRAY EQUIPMENT AND CONTAINERS.

DISCHARGE LIQUID WASTE IN PIT IN NON-CROP LANDS LOCATED AWAY FROM WATER SUPPLIES.

PERFORATE EMPTY METAL CONTAINERS.

CRUSH AND BURY GLASS.



÷ . '