(STAUFFER SHIELD)

IMIDAN® 50-WP GARDEN & HOME INSECTICIDE

A WETTABLE POWDER FOR CONTROL OF INSECTS ON DECIDUOUS SHADE TREES, ORNAMENTAL TREES, FRUIT TREES AND WOODY EYERGREENS

ACTIVE INGREDIENT:

N-(MERCAPTOMETHYL)P PHOSPHORODITHIOAT	HTH E)	IAI	[]	MII	DE	<u>S</u> -	- <u>((</u>	<u>), (</u>	<u>)</u> –1	11C	1E	THY	L •	ŧ	•		•	ı	•	50%
INERT INGREDIENTS:	•	•			•			ŧ		1	•	•			•	ı	5			50% 100%

U.S. PATENT NO. 2,767,194

WARNING

KEEP OUT OF REACH OF CHILDREN

ACCEPTED

SEP 2 6 1972

UNDER THE ALLEGAL INSECTICIDE ACT FOR EL COMIC POISON FEGISTER. ED UNDER NO. THE SUBJECT ACT OF THE SUBJECT

MAY BE HARMFUL IF SWALLOWED, IGHALED
OR ABSORBED THROUGH THE SKIN.

DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING.
DO NOT BREATHE DUST OR SPRAY MIST.
IN CASE OF CONTACT, WASH IMMEDIATELY
WITH SOAP AND WATER.
FLUSH EYES WITH WATER.
WASH HANDS AND FACE THOROUGHLY BEFORE EATING OR SMOKENG.
WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER
BEFORE RE-USE.
DO NOT CONTAMINATE FOOD OR FEED.

NOTE TO PHYSICIAN: IMIDAN IS AN ORGANOPHOSPHORUS INSECTICIDE. ATROPINE IS ANTIDOTAL.

EPA REG. NO

476-2112-AA

MADE IN U.S.A. BY

LBS. NET

STAUFFER CHEMICAL COMPANY WESTPORT, CONNECTICUT 06880

2-ARC-720802-12

PRODUCT INFORMATION

IMIDAN 50-WP is a wettable powder for use in water for spray applications. IMIDAN 50-WP is compatible with most commonly used insecticides and fungicides. It is not compatible with alkaline materials such as lime or bordeaux mixtures.

OBSERVE THE FOLLOWING USE PRECAUTIONS

- READ ALL LABEL DIRECTIONS BEFORE USING. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.
- . THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.
- , KEEP CHILDREN AND PETS AWAY FROM AREA WHILE SPRAYING.
- NEVER LEAVE INSECTICIDE PACKAGE, SPRAY MIXTURE OR SPRAYER UN-ATTENDED IF YOU ARE REQUIRED TO LEAVE AREA BEFORE SPRAYING IS COMPLETED.

STORAGE: STORE THIS PRODUCT IN A CABINET, STORAGE ROOM OR AREA THAT CAN BE LOCKED. NEVER STORE WITH OR NEAR FOOD, MEDICINES, OR HOUSEHOLD CLEANING MATERIALS. NEVER STORE EXCESS SPRAY (SEE "CLEANUP AND DISPOSAL").

CLEANUP AND DISPOSAL: WASH ALL EQUIPMENT (MEASURING SPOON, PAILS, SPRAYER, ETC.) INSIDE AND OUT WITH WATER. IMMEDIATELY DISCARD EXCESS SPRAY AND CLEANUP-WATER BY BURYING IN A SAFE PLACE LOCATED AWAY FROM WATER SUPPLIES. DISPOSE OF EMPTY CONTAINER BY BURYING WITH WASTES OR BY WRAPPING IN SEVERAL LAYERS OF NEWSPAPER AND PLACING IN TRASH FOR COLLECTION. IN CASE OF SPILLS, CLEAN UP IMMEDIATELY. WASH OFF WALKS AND OTHER PAVED AREAS BY HOSING.

FISH AND WILDLIFE CAUTIONS: THIS PRODUCT IS TOXIC TO FISH AND WILD-LIFE. DO NOT CONTAMINATE DOMESTIC OR IRRIGATION WATER SUPPLIES, LAKES, STREAMS OR PONDS BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED.

DIRECTIONS FOR USE

WHEN TO SPRAY: CHOOSE A COOL, CALM PERIOD, PREFERABLY IN EARLY MORNING OR EVENING. DO NOT APPLY IF RAIN IS EXPECTED OR BEFORE LEAF SURFACES ARE DRY.

EQUIPMENT: USE A COMPRESSED AIR SPRAYER, BUCKET PUMP SPRAYER, SLIDE PUMP SPRAYER, OR SMALL POWER SPRAYER OR WHEELBARROW TYPE SPRAYER. FOR SMALLER TREES AND SHRUBS, A HOSE PROPORTIONER DESIGNED FOR WETTABLE POWDERS CAN ALSO BE USED. FOR PEST CONTROL TREATMENT OF LARGE TREES, HIGH PRESSURE IS NEEDED TO DRIVE THE SPRAY INTO DENSE FOLIAGE AND TO REACH THE HIGHEST BRANCHES.

MIXING: PREPARE ONLY THE AMOUNT OF SPRAY NEEDED.

POUR RECOMMENDED AMOUNT (SEE RECOMMENDATIONS BELOW) OF THIS MATERIAL ON SURFACE OF WATER IN NEARLY-FILLED SPRAY TANK. ADD BALANCE OF WATER TO FILL TANK.

AGITATION: SPRAY MIXTURE REQUIRES AGITATION TO KEEP INSECTICIDE FROM SETTLING. WHEN USING SMALL EQUIPMENT, SHAKE SPRAYER BEFORE AND FREQUENTLY DURING APPLICATION. POWER EQUIPMENT: KEEP AGITATOR RUNNING DURING MIXING AND SPRAYING.

SPRAYING: SPRAY THOROUGHLY, COVERING LOWER AND UPPER LEAF SURFACES, STEMS, BRANCHES, TRUNK AND BASE OF THE PLANT UNTIL LIGHTLY DRIPPING.

RECOMMENDATIONS

USE 2 TABLESPOONFULS IMIDAN 50-WP PER GALLON OF WATER WHEN INSECTS OR THEIR DAMAGE APPEAR. REPEAT APPLICATION AS NECESSARY TO MAINTAIN INSECT CONTROL.

· · · · >

2= >-3=12 RECOMMENDED ON <u>DECIDUOUS FRUIT TREES</u> AS SPECIFIED BELOW (LIMIT USE TO 10 GALLONS SPRAY PER TREE PER APPLICATION):

PEST	APPLE	PEAR	PEACH	APRICOT/ NECTARINE	PRUNE/ PLUM	TART CHERRY
ELM SPANWORM* GYPSY MOTH* APHIDS (GREEN, ROSY) APPLE MAGGOT CODLING MOTH FRUIT TREE LEAFROLLER* PEAR PSYLLA PLUM CURCULIO RED BANDED LEAFROLLER TARNISHED PLANT BUG** ORIENTAL FRUIT MOTH PEACH TWIG BORER EUROPEAN RED MITES **	X X X X	X X X X	X	X X X	X	X X*
RED BANDED LEAFROLLER TARNISHED PLANT BUG** ORIENTAL FRUIT MOTH PEACH TWIG BORER EUROPEAN RED MITES **	X X	X	X X X	X X X	X X X	X

^{*}NORTHEASTERN U.S. ONLY.
**SUPPRESSION ONLY.

FOR BEARING TREES DO NOT PICK FRUIT WITHIN 7 DAYS OF SPRAYING (14 DAYS FOR PEACHES, APRICOTS AND NECTARINES).

RECOMMENDED ON <u>DECIDUOUS SHADE AND ORNAMENTAL TREES</u> (ASH, BEECH, OAK, DOGWOOD, WILLOW, HICKORY, HAWTHORNE, BIRCH, ELM, MAPLE) TO CONTROL:/

ELM SPANWORM / SPRING CANKERWORM / GYPSY MOTH /

. RECOMMENDED ON WOODY EVERGREENS (ARBORVITAE, CEDAR, FIR, HEMLOCK, JUNIPER, PINE, SPRUCE, YEW) TO CONTROL! -

. . 7

GYPSY MOTH

. RECOMMENDED ON BIRCH TREES TO CONTROL:

BIRCH LEAF MINERY

NOTE: LIMIT ALL USES FOR CONTROL OF GYPSY MOTH, ELM SPANWORM, SPRING CANKERWORM AND BIRCH LEAF MINER TO THE NORTHEASTERN U.S. ONLY.

NOTICE: STAUFFER CHEMICAL COMPANY MAKES NO WARRANTY, EXPRESS OF IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE, CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON STAUFFER'S LABEL.