



# CAPTAN-DIAZINON-LINDANE<sup>\*</sup> 33.5-11-16.6 SEED PROTECTANT

PLANTER BOX APPLICATION FOR PROTECTION OF SEEDS AND SEEDLINGS

SPECIAL LUBRICANT ADDED—DISCOLORED

ACCEPTED  
SEP 19 1973  
UNION PACIFIC FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON PATENT  
STAUFFER NO. 476-2115

**ACTIVE INGREDIENTS:**

CAPTAN: N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide . . . . .	33.5%
O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl)phosphorothioate* . . . . .	11.0%
Gamma isomer of benzene hexachloride (from Lindane) . . . . .	16.6%
<b>INERT INGREDIENTS:</b> . . . . .	38.9%
	100.0%

\*DIAZINON, a Registered Trademark of Ciba-Geigy Corporation.

**OBSERVE THE FOLLOWING USE PRECAUTIONS**

Read all directions before using and use only as specified on this label.  
 Do not use on seed already treated with an insecticide except for Malathion or Methoxychlor used for control of stored grain insects.  
**Fish and Wildlife Cautions:** Toxic to fish and wildlife, keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Treated seed on the soil surface will be hazardous to birds and other wildlife. Dispose of excess treated seed by burial.  
**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED.  
DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.

The federal government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed. Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

**RECOMMENDATIONS**

CAPTAN-DIAZINON-LINDANE 33.5-11-16.6 SEED PROTECTANT is a special formulation for the planter box treatment of beans, corn, peas, sorghum and soybeans for the protection of seeds against seed corn beetles, seed corn maggots, wireworms, damping-off, seed rots, seedling blights and fungus diseases of seeds and seedlings.

**DIRECTIONS FOR USE**

**Beans, Corn, Peas, Sorghum, Soybeans — Planter Box Application:** Use 3 ounces per bushel of seed. Fill the drill box one half full of seed then treat with one half the amount of seed protector and mix well by stirring with a stick or paddle until color is uniform in the seed. Add the remaining seed to drill box and cover with the remainder of the seed protectant. Mix thoroughly as above. Regularly inspect planter plates and clean as required. If the seed treatment increases the friction between seeds, sufficient to reduce the rate of delivery through a grain drill, add 1 ounce of graphite per bushel as a drill lubricant. Adjust the seed drill to give the required rate of seed delivery with treated seed.

**12 OUNCE CONTAINER — TREATS 3 BUSHELS.**

**WARNING** Keep Out of Reach of Children  
 MAY BE HARMFUL OR FATAL IF SWALLOWED.  
 MAY BE ABSORBED THROUGH THE SKIN.  
 Do not breathe dust.  
 Do not get in eyes, on skin or on clothing.  
 Wash thoroughly with soap and water after handling.  
 Do not contaminate food or feed.

PROTECT FROM EXCESSIVE HEAT • STORE IN A COOL DRY PLACE

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES

*NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.*

EPA Reg. No. 476-2115-AA

ONE OUNCE MEASURING CUP ENCLOSED.

NET WEIGHT 12 OUNCES

Made In U.S.A. By

**STAUFFER CHEMICAL COMPANY**

WESTPORT, CT 06880