


# TRITHION-SULFUR 3-50 DUST

AGRICULTURAL INSECTICIDE-ACARICIDE & FUNGICIDE



**ACTIVE INGREDIENTS:**

Carbophenothion: S-[(p-chlorophenylthio)methyl] O,O-diethyl phosphorodithioate	3%
Sulfur	50%
<b>INERT INGREDIENTS:</b>	<b>47%</b>
	<b>100%</b>

U. S. Pat. No. 2,793,224

E.P.A. Reg. No. 476-2118

**CAUTION** **Keep Out of Reach of Children**

**HAZARDOUS IF SWALLOWED. AVOID BREATHING DUST, CONTACT WITH SKIN AND EYES.** Wear dust respirator and goggles. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate food and feed. Wash all contaminated clothing with soap and hot water before re-use.

**ANTIDOTE**—TRITHION may cause cholinesterase inhibition. **Atropine** is antidotal. **2-PAM** is also antidotal when administered early, and in conjunction with atropine.

**USE PRECAUTIONS**

**READ ALL LABEL DIRECTIONS BEFORE USING.**

Use only on crops listed and only as specified on this label.

This produce is toxic to bees and should not be applied when bees are actively visiting the area.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damages caused by drift from his property to that of others.

SULFUR may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply SULFUR to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

SULFUR will cause severe fruit and leaf injury to SULFUR-sensitive crops. Do not apply or allow to drift to apricots, anjou or comice pears, cranberries, cucurbits (cucumber, cantaloupes, melons, squash), walnuts, or other sensitive plants.

When growing crops for processing, consult the processor before applying SULFUR.

Consult State Agricultural Experiment Station or the Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.

**FISH AND WILDLIFE CAUTIONS:** Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Birds feeding on treated areas may be killed.

50 LB. NET

**DIRECTIONS FOR USE**

Apply the specified amount of this product uniformly to the foliage when all is calm, using ground equipment or aircraft. Apply at the first sign of infestation and infection. Repeat at regular intervals when needed to maintain control, but observe use limits for specific crops.

For control of diseases after the prescribed time limits have expired, an effective fungicide should be used alone, in a manner which will not leave residues in excess of established tolerances.

**RECOMMENDATIONS**

Maximum permissible rate per acre, expressed as TRITHION-SULFUR 3-50 DUST, is given in parenthesis ( ), after each crop claim.

**ALFALFA, CLOVER:** Meadow Spittlebug, Pea Aphid, Potato Leafhopper, Spotted Alfalfa Aphid (alfalfa only), Two-Spotted Mite—Apply 20 to 33 lb. per acre. Apply 33 lb. per acre for the control of the meadow spittlebug. Do not apply more than one application per cutting. Do not graze or cut hay within 28 days after application. (33 lb.)

**BEANS, SNAP:** Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Powdery Mildew, Serpentine Leaf Miner, Two-Spotted Mite—Apply 20 lb. per acre. Do not apply within 7 days of harvest. Do not apply more than two applications per season. Do not graze or cut hay within 21 days after application. (20 lb.)

**BEANS, DRY (SHELLED):** Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Powdery Mildew, Serpentine Leaf Miner, Two-Spotted Mite—Apply 20 lb. per acre. Do not apply more than two applications per season. Do not graze or cut hay within 21 days after application. (20 lb.)

**CHERRY, PEACH, PLUM, PRUNE:** Aphids, Orchard Mites, Powdery Mildew—Apply 30-40 lb. per acre. Do not apply within 30 days of harvest. (60 lb.)

**COTTON:** Carmine (Cinnabar) Mite, Cotton Leafworm, Tropical Mite, Two-Spotted Mite—Use 20 lb. per acre. If mite populations are high, use 20-33 lb. per acre and repeat applications in 5-10 days. Aphids, Cotton Leaf Perforator, Fleahopper—Use 33 lb. per acre. Treated plants should not be grazed or fed to livestock. (33 lb.)

**GRAPES:** Grape Leafhopper, Powdery Mildew, Two-Spotted Mite—Apply 25-33 lb. per acre. Do not apply within 30 days of harvest. Do not apply to sensitive varieties such as: Charbono, Gamay, Golden Muscat, Muscat and Italia. (33 lb.)

**PEA:** Pea Aphid, Powdery Mildew, Serpentine Leaf Miner, Two-Spotted Mite—Apply 20-25 lb. per acre. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock. (40 lb.)

**SEED CROPS (Alfalfa, Clover, Sugar Beets and Vegetable Crops listed on this label):** Bean Aphid, Brown Wheat Mite, Pea Aphid, Serpentine Leaf Miner, Spotted Alfalfa Aphid, Two-Spotted Mite—Apply 30 lb. per acre. Do not graze or feed crop residue to livestock. Sugar beet seed crops can be treated up to 14 days before harvest and crop residue can be fed to livestock.

**SUGAR BEETS:** Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two-Spotted Mite—Apply 20-30 lb. per acre. Do not apply within 14 days of harvest. (33 lb.)

**STRAWBERRY:** Powdery Mildew, Strawberry Aphid, Two-Spotted Mite—Apply 20-30 lbs. per acre. Do not apply within 3 days of harvest. (40 lb.)

**TOMATO:** Potato Flea Beetle, Potato Leafhopper, Potato Psyllid, Serpentine Leaf Miner, Tomato Russet Mite, Tropical Mite, Two-Spotted Mite—Apply 20-30 lb. per acre. Do not apply within 7 days of harvest. (40 lb.)

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label. ARC-720306-0

BURY WASTE AND EMPTY CONTAINER IN A SAFE PLACE

Made in U.S.A. by

**STAUFFER CHEMICAL COMPANY**

WESTPORT, CT 06880