



# Dyfonate<sup>®</sup> 5-G

## GRANULAR ORNAMENTAL TURF INSECTICIDE

(FOR USE BY COMMERCIAL APPLICATORS AND TURF MANAGEMENT PERSONNEL IN THE SOUTHEASTERN U.S.)

### ACTIVE INGREDIENT:

O-ethyl S-phenyl ethylphosphonodithioate (U.S. Pat. No. 2,988,474)..... 5%

### INERT INGREDIENTS:

..... 95%  
100%

Contains 5 pounds active ingredient per 100 pounds.

## WARNING

Keep Out of Reach  
of Children

### HARMFUL OR FATAL IF SWALLOWED.

Do not get on skin, in eyes, on clothing.

Avoid inhaling dust and skin contact.

Wear gloves when handling.

In case of contact, wash immediately with soap and water.

Always wash thoroughly after handling.

**NOTE TO PHYSICIANS:** Exposure may cause cholinesterase inhibition. **ATROPINE IS ANTIDOTAL. 2-PAM** also is antidotal when administered early, and in conjunction with atropine.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

EPA Reg. No. 476-2121

2-ARC-720919-4

### USE PRECAUTIONS

- Read all directions before using.
- Do not apply in any manner not specified on this label.
- For use by commercial applicators and turf management personnel only.
- Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops.
- Keep people and pets out of treated areas until grass is dry.
- This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**DISPOSAL:** COMPLETELY EMPTY THE CONTENTS OF BAG AND BURY UNUSED CHEMICAL AT LEAST 18 INCHES DEEP IN AN ISOLATED LOCATION AWAY FROM WATER SUPPLIES. BURN EMPTY BAG IMMEDIATELY.

### RECOMMENDATIONS AND DIRECTIONS FOR USE

For control of Chinch Bug in established ornamental turf in Southeastern United States.

DO NOT PREWATER THE TURF AREA.

Apply at first sign of insects or insect damage. Use 60 to 80 pounds per acre (1½ to 1¾ pounds per 1,000 square feet) applied uniformly with a granular pesticide applicator. Immediately after application, water the treated turf thoroughly for 10 to 15 minutes to wash the product into the root zone. Retreat as needed to keep local infestations under control.

50 LB. NET

# STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

476-2121

10-04-1972

1/2

DATA  
SEARCH

ACCEPTED  
OCT 04 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. .... SUBJECT  
TO ATTACHED COMMENTS.

476-2121

2/6