DVFORTER 2-G A GRANULAR SOIL INSECTICIDE FOR LAWNS AND HOME GARDENS

ACTIVE INGREDIENT:

O-ethyl S-phenyl ethylphosphonodithioate
INERT INGREDIENTS:

% BY WGT. 2% 98%

100%

Ummula

U.S. Patent No. 2,988,474

CAUTION

HARMFUL IF SWALLOWED.

Do not get on skin, in eyes, on clothing. Avoid inhaling dust and skin contact.

Wear gloves when handling.

Always wash thoroughly after handling.

Keep Out of Reach of Children?

ACCEPTED

AUG 5 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Acts as amended, for the pesticide registered under EPA Reg. No. 4724 2726

NOTE TO PHYSICIANS: Exposure may cause cholinesterase inhibition. ATRO-PINE IS ANTIDOTAL. 2-PAM also is antidotal when administered early, and in conjunction with atropine.

USE PRECAUTIONS

- Read all directions before using.
- Wash any spills off walks by hosing.
- Avoid contamination of food and feed.

FISH AND WILDLIFE CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

STORAGE — Store this product in a locked cabinet away from living quarters.

THIS PRODUCT CONTROLS

Chinch Bugs Earwigs Sod Webworms Garden Symphylans (Centipedes) Wireworms Cabbage Maggots

DIRECTIONS FOR USE

LAWNS — Use on established Bahia, Bermudagrass, Bluegrass, St. Augustine, Zoysia. Do not prewater the lawn. Apply at the first sign of insects or insect damage.

For the control of CHINCH BUGS. EARWIGS and SOD WEBWORMS use $3\frac{1}{2}$ - $4\frac{2}{3}$ lbs. per 1,000 sq. ft. Use the higher rate for heavy infestations. Recommended for Chinch Bugs throughout the Southern U.S.; for Earwigs and Sod Webworms in the southeast.

Apply uniformly using a lawn fertilizer spreader. Immediately after application water the treated area thoroughly for 10-15 minutes to wash granules into the root zone.

Repeat as necessary to keep infestations under control. Do not allow children or pets on treated area until completely dry.

GARDENS (Recommended Nationally)

Use DYFONATE 2-G only on the following vegetables at the rate of application shown.

	Use these amounts per 100 sq. ft. to control			
Vegetable	Garden Symphylans	Wireworms	Cabbage Maggot	Remarks
Beans, Snap	1/4 lb.	½ lb.		
Beans, Lima	⅓ lb.			
Beets	1/4 lb.		¹ / ₄ lb.	
Broccoli	1/4 lb.			
Brussels Sprouts	1/4 lb.			Apply
Cabbage	⅓ lb.			just
Cauliflower	¹∕a lb.			prior to
Corn	1/4 lb.	1/2 lb.		planting
Potatoes-White	⅓ lb.	¹ / ₂ lb.		or
Potatoes-Sweet		1/2 lb.		transplanting.
Radishes	½ lb.		≟a lb.	
Tomatoes	1.4 lb.			
Strawberries	⅓ lb.			

HOW TO APPLY — Shake from canister or spread with fertilizer spreader uniformly before planting or at the time of planting. Mix into the top 2-3 inches of soil with a hand rake or power tiller.

DISPOSAL — Do not re-use container. Puncture, wrap in several layers of newspaper and place in trash for collection.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

EPA Reg. No. 476-2120

NET WGT.

A-2

Made in U.S.A. By

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

Stauffer GRANULAR ORNAMENTAL TURF INSECTICIDE

FOR USE BY COMMERCIAL APPLICATORS AND TURF MANAGEMENT PERSONNEL

ACTIVE INGREDIENT:

% by wgt.

O-ethyl S-phenyl ethylphosphonodithioate

INERT INGREDIENTS:

95% 100%

Contains 5 pounds active ingredient per 100 pounds.

(U.S. Pat. No. 2.988,474)

WARNING

of Children

HARMFUL OR FATAL IF SWALLOWED.

Do not get on skin, in eyes, on clothing.

Avoid inhaling dust and skin contact.

Wear gloves when handling.

In case of contact, wash immediately with soap and water.

Always wash thoroughly after handling.

NOTE TO PHYSICIANS: Exposure may cause cholinesterase inhibition. ATRO-PINE IS ANTIDOTAL. 2-PAM also is antidotal when administered early, and in conjunction with atropine.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

EPA Reg. No. 476-2121

Keep Out of Reach

RECOMMENDATIONS

For use on established turf: Bahia

Bermudagrass Bluegrass

For control of:

Chinch Bugs

Earwigs Recommended for Chinch Bugs in the Southern U.S.; on Earwigs and Sod Webworms in the Southeastern U.S.

DIRECTIONS FOR USE Do not prewater the turf area.

Apply at first sign of insects or insect damage.

Use 60-80 lbs. DYFONATE 5.G per acre (1½ to 1¾ lbs. per 1,000 sq. ft.) Apply uniformly using a granular pesticide applicator. Immediately after application, water the treated area for 10-15 minutes to wash granules into the root zone.

Repeat as needed to keep local infestations under control. Do not allow people or pets on treated area until grass is dry.

50 PCUNDS NET (22.7 KG.)

Made in U.S.4 By

USE PRECAUTIONS

- Read all directions before using.
- Apply only as directed on label.
- Do not allow to drift onto adjoining food, fiber or pasture crops.
- This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment. or disposal of wastes

STORAGE — Store unused product in locked area away from food or feed crops. **DISPOSAL** — Make sure bag is completely empty. Bury waste and empty bag* ... in non-crop area away from water supplies.

St. Augustine

Sod Webworms

Zoysia

ACCEPTED

AUG 5 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act. as amended, for the pesticide registered under EPA Reg. No.

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880