

# How to use SANITIZER on the farm

- **Clenesco SANITIZER** is both a cleaner and a sanitizer. It reduces bacteria counts and reduces your cleaning time.
- The standard **Clenesco SANITIZER** solution containing 200 p.p.m. active sanitizing ingredient is prepared with one ounce of **Clenesco SANITIZER** in each 7½ quarts of water. The scoop in this container, level-full of **Clenesco SANITIZER** is sufficient for 10 quarts of water.

## MILKING MACHINES AND HOSES

1. Just before milking, re-assemble cleaned milker and draw standard **Clenesco SANITIZER** solution through it, allowing contact time of two minutes at 75 F. or higher. Allow equipment to drain prior to beginning of milking operation.
2. Save ¼ of this solution for rinsing equipment after milking.
3. Immediately after milking, draw clear water through milkers to remove milk film. Continue until water runs clear.
4. Add warm water to the ¼ used solution from step #2 and draw this solution through milkers to remove residual milk.
5. Dismantle and wash with brush in standard **Clenesco SANITIZER** solution at 110-115 F.
6. Rinse in second tank, using clear water. Store dry to protect all milk-contact surfaces.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

LOT NO.

ACCEPTED

7-13-73  
476-2124



CLENESCO®

# SANITIZER

EPA Reg. No. 476-2124

CLEANER-SANITIZER FOR  
DAIRY FARM UTENSILS

### ACTIVE INGREDIENTS:

Sodium Carbonate .....	26.60%
n-Alkyl (C <sub>12</sub> -50%, C <sub>14</sub> -30%, C <sub>16</sub> -17%, C <sub>18</sub> -3%)	
dimethyl benzyl ammonium chloride.....	5.00%

<b>INERT INGREDIENTS*</b> .....	68.40%
---------------------------------	--------

\*Active as cleaning agents.

ARC-730305

**DANGER**

Keep Out of Reach  
of Children

**CAUSES SEVERE BURNS.**

**HARMFUL IF SWALLOWED.**

Do not get in eyes, on skin, on clothing.

Avoid breathing dust.

Protect eyes and skin when handling.

Avoid contamination of food and feed.

**FIRST AID** IN CASE OF CONTACT, IMMEDIATELY flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

(See right panel for internal antidote.)

**NET WEIGHT — 25 POUNDS**

**STAUFFER CHEMICAL COMPANY**  
**INDUSTRIAL CHEMICAL DIVISION**  
WESTPORT, CT 06880

**INTERNAL ANTIDOTE:** If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

## FARM UTENSILS— PAILS, STRAINERS

1. Rinse with water immediately after use.
2. Dismantle if assembled. Brush all surfaces with warm **Clenesco SANITIZER** solution. Rinse with clear water.
3. Store to protect all milk-contact surfaces.
4. Before milking, rinse all surfaces with a fresh 200 p.p.m. **Clenesco SANITIZER** solution at 75 F. or above permitting two minutes contact time. Allow equipment to drain prior to beginning milking operation.

Complete directions for cleaning and sanitizing other dairy farm equipment are available from your dairy. The regulations of your fieldmen and of local health authorities should always be followed. Consult your dairy when you need help. Do not mix with soap or other cleaners. Use a freshly made solution for each milking.

## COWS' UDDERS

1. Saturate a large clean cloth or paper towel with warm standard **Clenesco SANITIZER** solution. Use this saturated cloth to wash udders and teats.
2. Wring cloth dry. Use it to wipe excess solution off udders.
3. Use a clean towel for each animal. Never re-use towel or dip used towel back into solution. Always use freshly prepared solution and non-corrodible utensils.

**DISPOSAL:** Do not reuse empty container. Discard it.

A standard use solution of **Clenesco SANITIZER** containing 200 parts per million active ingredient is at a pH higher than 6.0 and contains more than 0.2% compatible complex phosphate. This solution has a 99.999% effective count reduction with a 30 second exposure in waters up to 500 p.p.m. hardness. This product fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance—1965 Recommendations of the U. S. Public Health Service in waters up to 500 p.p.m. of hardness, calculated as CaCO<sub>3</sub> when tested by the A.O.A.C. Germicidal and Detergent Sanitizers—Official Method.