



# TOXAPHENE 40-WP

## AGRICULTURAL INSECTICIDE

### WETTABLE POWDER

**ACTIVE INGREDIENT:**

Toxaphene (Technical Chlorinated Camphene)\* 40%

**INERT INGREDIENTS:**

60%  
100%

\*Chlorine content 67% to 69%.

**WARNING**

**Keep out of Reach of Children**

**MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN.** Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not contaminate food or feed.

**OBSERVE THE FOLLOWING USE PRECAUTIONS**

READ ALL LABEL DIRECTIONS BEFORE USING.

Use only for crops listed and only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

In order that pesticide residues on food and forage crops will not exceed tolerances established by the Environmental Protection Agency, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crop as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the timing of applications needed will vary with the local conditions.

**FISH AND WILDLIFE CAUTIONS:** Highly toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate any water supply or apply under conditions favoring drift outside the treated area.

Rinse spray equipment. Bury liquid waste in pit on non-crop lands located away from water supplies.

**BURY WASTE AND EMPTY CONTAINER IN A SAFE PLACE.**

**4 LBS. NET**

EPA Reg. No. 476-2133

ARC 721031/0

Made in U.S.A. by

**STAUFFER CHEMICAL COMPANY**

**WESTPORT, CT 06880**

TOXAPHENE 40-WP

**DIRECTIONS FOR USE**

Measure the specified amount of the product and mix with water in a clean, preferably lined, tank or premix in a clean, preferably lined, tank. Add 100 parts of water to 1 part of product. Keep agitator running during filling and pouring operations. Do not use if the agitator is not started. Do not use in low volume gear pump spray equipment. Do not combine with incompatible liquids unless previous use of the mixture has been previously tested and found safe to plants. Emulsifiable liquids should always be thoroughly mixed with product in a tank of water before adding this product.

**RECOMMENDATIONS**

Dosage rates are given in terms of Toxaphene (40% WP) per acre in sufficient water to provide thorough coverage (usually 7 to 10 gallons water per acre by hand, 20 to 25 gallons water per acre by ground equipment). Except where otherwise noted, all applications at the first sign of infestation and repeat as needed to maintain control but observe use limitations specified for certain crops.

Maximum permissible rate per acre, expressed as TOXAPHENE 40-WP, given in parenthesis, ( ) after each crop claim.

**BEANS:** Armyworm, Corn Earworm, Cutworms (Climbing and Surface-feeding), Thrips — Use 5 pounds. Bean Leaf Beetle, Blister Beetle, Cowpea Curculio, Cucumber Beetle, Flea Beetle, Grasshoppers, Lygus Bug, Mexican Bean Beetle, Wax Bean Beetle, Serpentine Leaf Miner — Use 5 to 10 pounds. Bean Leafhopper, Bean Leaf Roller — Use 2½ pounds. Bean Leaf Skeletonizer, Garden Webworm, Pea Weevil — Use 7½ to 10 pounds. Salt Marsh Caterpillar — Use 10 pounds. Do not apply to Green or Snap Beans after pods start to form. Do not feed treated vines to dairy animals or animals being finished for slaughter. (15 pounds)

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Imported Cabbageworm, Fall Armyworm, Green Peach Aphid, Serpentine Leaf Miner, Thrips — Use 3½ to 5 pounds. Cabbage Looper, Corn Earworm, Diamond-back Moth Larvae — Use 3¾ to 10 pounds. Cutworms (Climbing and Surface-feeding) — Use 5 pounds. Flea Beetle — Use 7½ pounds. Do not apply after heads or edible parts begin to form or within 30 days of harvest on Broccoli, Brussels Sprouts and Cauliflower. Do not apply after heads start to form on Cabbage. (Cabbage — 10 pounds, all others — 20 pounds.)

**CELERY:** Imported Cabbageworm, Fall Armyworm, Green Peach Aphid, Serpentine Leaf Miner, Thrips — Use 3¾ to 5 pounds. Celery Leaf Tier, Vegetable Weevil — Use 10 to 12½ pounds. Cutworms (Climbing and Surface-feeding) — Use 5 to 12½ pounds. Do not apply after celery is half grown or starts to bunch. (12½ pounds)

**CORN:** Billbug — Use 5 pounds. Budworm (Fall Armyworm and Corn Earworm feeding in whorl), Cutworms, Flea Beetle, Grasshopper — Use 5 to 7½ pounds. Corn Lantern Fly — Use 3¾ to 10 pounds. Lesser Corn Stalk Borer — Use 10 pounds. Whiteworm — Use 5 to 6½ pounds. Do not feed forage to dairy animals or animals being finished for slaughter. (15 pounds)

**COTTON:** Armyworms, Boll Weevil, Bollworm, Cotton Leafworm, Cutworms, Darkling Beetle, Grasshopper, Plant Bugs (Tarnished, Rapid and Lygus) — Use 5 to 10 pounds. Flea Beetle, Leafhopper, Thrips — Use 2 to 7½ pounds. Garden Webworm — Use 7½ pounds. Cotton Leaf Perforator — Use 10 pounds. Do not graze dairy animals or animals being finished for slaughter in Cotton fields treated late in season. Do not feed gin waste to livestock. After fields are open applications at 10 pound rate should not be made more frequently than every 5 days. (10 pounds)

**COWPEA (INCLUDING FIELD PEAS):** Bean Leafhopper, Bean Leafroller — Use 2½ pounds. Cowpea Curculio, Lygus Bug — Use 5 to 7½ pounds. Cutworms, Darkling Ground Beetle — Use 7½ pounds. Do not apply to varieties with edible pods after pods start to form. Do not feed treated forage to dairy animals or animals being finished for slaughter. Where Peas are taken from pod and fed separately, do not apply within 7 days of harvest. (7½ pounds)

**EGGPLANT:** Cutworms (Climbing and Surface) — Use 5 pounds. Cucumber Beetle — Use 7½ pounds. Armyworm, Blister Beetle, Colorado Potato Beetle, Flea Beetle, Leafhopper, Serpentine Leaf Miner — Use 5 to 7½ pounds. Do not apply within 5 days of harvest. (7½ pounds)

**LETTUCE:** Armyworms, Cutworms — Use 5 pounds. Imported Cabbageworm, Fall Armyworm, Flea Beetle — Use 3¾ to 7½ pounds. Field Cricket, Grasshopper — Use 5 to 10 pounds. Cabbage Looper, Salt Marsh Caterpillar — Use 7½ to 12½ pounds. Do not apply after heads start to form. Do not apply to Head Lettuce after heads are on the table. (12½ pounds)

**PEPPERS:** Armyworms, Blister Beetle, Colorado Potato Beetle, Cutworms, Flea Beetle, Grasshopper, Serpentine Leaf Miner — Use 5 to 7½ pounds. Field Cricket, Pepper Weevil — Use 7½ pounds. Do not apply within 5 days of harvest. (7½ pounds)

**TOMATOES:** Cabbage Looper, Cutworms, Flea Beetle, Leafhopper, Lygus Bug, Salt Marsh Caterpillar, Armyworms, Blister Beetle, Colorado Potato Beetle, Field Cricket, Flea Beetle, Grasshopper, Imported Cabbageworm, Tomato Fruitworm — Use 5 to 10 pounds. Thrips — Use 5 to 7½ pounds. Do not apply within 3 days of harvest. (12½ pounds)

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Made in U.S.A. by

Stauffer Chemical Company, Westport, Connecticut