.....

MIDAN 12.5-WP ACCEPTED

GARDEN AND HOME INSECTICIDE

A WETTARLE POWDER FOR CONTROL OF INSECTS ON DECIDUOUS FRUIT TREES. SHADE **ORNAMENTAL TREES AND WOODY EVERGREENS**

ACTIVE INGREDIENT:

N-(mercaptomethyl)phthalimide S-(O,O-dimethyl phosphorodithioate) 12.5% 87.5% INERT INGREDIENTS: 100.0%

U.S. Patent No. 2,767,194

CAUTION

Keep Out of Reach of Children

MAY BE HARMFUL IF SWALLOWED. Avoid contact with skin, eyes or clothing. Avoid breathing of dust or spray mist. In case of contact, wash immediately with soap and water. Flush eyes with water. Wash hands and face thoroughly before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not contaminate food or feed.

NOTE TO PHYSICIAN: IMIDAN is an organophosphorus insecticide. ATROPINE IS ANTIDOTAL.

IMIDAN 12.5-WP is a wettable powder for use in water for spray applications. IMIDAN 12.5-WP is compatible with most commonly used insecticides and fungicides. It is not compatible with alkaline materials such as lime or Bordeaux mixtures.

OBSERVE THE FOLLOWING USE PRECAUTIONS

- . READ ALL LABEL DIRECTIONS BEFORE USING, Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative agricultural extension service. Keep children and pets away from area while spraying.

 Never leave insecticide package, spray mixture or sprayer unattended if you are required to leave
- area before spraying is completed.

STORAGE: Store this product in a cabinet, storage room or area that can be locked. NEVER store with or near food, medicines, or household cleaning materials. NEVER store excess spray (see "Cleanup and

CLEANUP AND DISPOSAL: Wash all equipment (measuring spoon, pails, sprayer, etc.) inside and out with water. Immediately discard excess spray and cleanup-water by burying in a safe place located away from water supplies. Dispose of empty container by burying with wastes or by wrapping in several layers of newspaper and placing in trash for collection. In case of spills, clean up immediately. Wash off walks and other paved areas by hosing.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

WHEN TO SPRAY: Choose a cool, calm period, preferably in early morning or evening. Do not apply if rain is expected or before leaf surfaces are dry.

EQUIPMENT: Use a compressed air sprayer, bucket pump sprayer, slide pump sprayer, or small power sprayer or wheelbarrow type sprayer. For smaller trees and shrubs, a hose-end sprayer designed for wettable powders can also be used. For pest control treatment of large trees, high pressure is needed to drive the spray into dense foliage and to reach the highest branches.

MIXING: Prepare only the amount of spray needed.

Pour recommended amount (see RECOMMENDATIONS below) of this material on surface of water in nearly-filled spray tank. Add balance of water to fill tank. For hose-end sprayers follow manufacturer's directions for using with wettable powders.

AGITATION: Spray mixture requires agitation to keep insecticide from settling. When using small equipment, shake sprayer before and frequently during application. POWER EQUIPMENT: Keep agitator running during mixing and spraying.

SPRAYING: Spray thoroughly, covering lower and upper leaf surfaces, stems, branches, trunk and base of the plant until lightly dripping.

RECOMMENDATIONS

EPA Reg. No. Use 3 tablespoonfuls IMIDAN 12.5-WP per gallon of water when insects or their damage appear. Repeat application as necessary to maintain insect control.

 Recommended on DECIDUOUS FRUIT TREES as specified below. Spray all sides of leaves and fruit thoroughly. Large trees may require up to 10 gallons of spray per tree.

PEST	Apple	Pear	Peach	Nectarine	Plum	Cherry	Grape
Cherry Fruit Fly*						X	
Elm Spanworm*	. X	X					
Grape Berrymoth*							Х
Grape Leathopper†*				_			X
Grape Leaffolder							Х
Gyosy Moth*		Х					
Aphids (Green Apple)	. x	X					
Aphids (Rosy Apple)		X					
Apple Maggot				x	Х		
Codling Moth		X					
Fruit Tree Leafroller*	Ŷ	Ÿ				X	
Contiverous Leafroller							X
Pear Psylla		x					
Plum Curculio	Y	Ŷ	Y	Y	X	X*	
Ped Banded Leafroiler		Ŷ	^	Ŷ	Ŷ	-1	
Tarnished Plant Bug**		Ŷ	Y	^	^		
Oriental Fruit Moth	^	•	Ŷ	Y	Y		
Oriental Fruit Moth			÷	•	Ŷ	v	
Peach Twig Borer		~	\$	•	Ş	Ç.	
European Red Mites**		\$	•	\$	\$	Ç	
Two-Spotted Mites**	^	^		^	^	^	

^{*}Northeastern U.S. only.

For bearing trees do not pick fruit within 7 days of spraying (14 days for Peaches, Apricots, Nectarines

APPLE MAGGOT, CODEING MOTH, ORIENTAL FRUIT MOTH, PLUM CURCULIO: Apply at petal fall and every 10 to 14 days thereafter until control is achieved, or consult your cooperative extension service for timing of applications.

- RECOMMENDED ON DECIDUOUS SHADE AND ORNAMENTAL TREES (ash, beech, oak, dogwood, wiflow, locust, bickory, hawthorn, birch, elm, maple) to control: ELM SPANWORM, SPRING CANKERWORM,
- RECOMMENDED ON WOODY EVERGREENS (arborvitae, cedar, fir, hemlock, juniper, pine, spruce, yew) to control: GYPSY MOTH.
- RECOMMENDED ON BIRCH TREES to control: BIRCH LEAF MINER.

NOTE: Limit all uses for control of Gypsy Moth, Elm Spanworm and Birch Leaf Miner to the Northeastern U.S. only. Recommended for Spring Cankerworm in the Northeastern and Midwestern U.S.

NOTICE: Stauffer Chemical Company makes no warranty, express or Implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

EPA Reg. No. 476-2161

A-1

registered under 476-216

LBS. NET

STAUFFER CHEMICAL COMPANY

^{**}Suppression only

tapply when most nymphs hatch, or consult your cooperative extension service for timing of applications.