

BEST DOCUMENT AVAILABLE

GENERAL USE PRECAUTIONS:

Read the entire label before using. Use only as recommended for claims listed.

During commercial exposure in seed treatment and loading operations, wear goggles, clean synthetic rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR part 11.

Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

SPECIAL USE PRECAUTIONS:

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED

NOT USE FOR FEED, FOOD OR OIL PURPOSES

The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

Dispose of excess treated seed by burial.

ENVIRONMENTAL HAZARD:

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Bury wastes and empty container in isolated area away from water supplies.

STATEMENT OF PRACTICAL TREATMENT:

IN CASE OF EYE CONTACT: Immediately flush eyes with running water for at least 15 minutes. Call a physician.

IN CASE OF SKIN CONTACT: Wash the affected area with soap and warm water.

PRODUCT INFORMATION:

CAPTAN 4 FLOWABLE SPR is a liquid suspension especially formulated for the treatment of seed prior to storage and planting to protect seed from molds and other fungi causing storage loss, and to protect seed from seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights.

DIRECTIONS FOR USE AND RECOMMENDATIONS:

It is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions before using and use only as specified on this label.

IMPORTANT: Before using - stir thoroughly to mix contents.

DIRECTIONS: Thoroughly mix the recommended amount of CAPTAN 4 FLOWABLE SPR into the required amount of water for the slurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations.

BEST DOCUMENT AVAILABLE

ACCEPTED

JUN 22 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, the pesticide NO. 471-2192



CAPTAN

4 FLOWABLE SPR SEED PROTECTANT — Dye Added Agricultural Fungicide — Liquid Suspension

ACTIVE INGREDIENT:

CAPTAN: N[(Trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

39.0%

INERT INGREDIENTS:

61.0%

100.0%

Contains 4 Pounds CAPTAN Per Gallon

CAUTION — Keep Out of Reach of Children

- Causes eye and skin irritation.
- Harmful if swallowed, inhaled or absorbed through skin.
- Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling.
- Avoid inhaling spray mist.
- Do not use or store near food or feed.

See side panel for additional precautionary statements and statement of practical treatment.

STORE IN A COOL, DRY PLACE.
STIR THOROUGHLY TO MIX CONTENTS BEFORE USING.

5 GAL. NET

Made in U.S.A. by

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

SEED TREATMENT DOSAGE

SEED

- Alfalfa, clover, lespedeza, trefoil
- Beans (snap, dry, cowpeas)
- Beets (garden)
- Cereal grains (for seed and seedling rots not for bunt smut)
 - Barley, rye
 - Oats
 - Wheat
- Cole crops (broccoli, cabbag, rutabaga)
- Corn
 - Field
 - Sweet
- Cotton - Acid delint, Reginned fuzzy
- Cucumber, cantaloupe
- Crucifers (Mustard, radish, rape, turnips)
- Flax
- Grass
- Milo
- Peanuts (unshelled)
- Peas, Lentils
- Peppers
- Rice
- Sorghum (hulled) - For Kernel smut
- Soybeans
- Spinach
- Squash, pumpkin, watermelon, muskmelon
- Sugar beets - West, East
- Sunflowers
- Swiss chard

Further specific information on the subject of seed treatment should be obtained from State Agricultural Experiment Stations or Agricultural Extension Service.

BEST DOCUMENT AVAILABLE

A

B

ACCEPTED
 JUN 29 1979
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, No. 47-2192



CAPTAN

4 FLOWABLE SPR

SEED PROTECTANT — Dye Added

Agricultural Fungicide — Liquid Suspension

ACTIVE INGREDIENT:
 CAPTAN: N[(Trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide 39.0%

INERT INGREDIENTS: 61.0%

Contains 4 Pounds CAPTAN Per Gallon 100.0%

CAUTION — Keep Out of Reach of Children

- Causes eye and skin irritation.
- Harmful if swallowed, inhaled or absorbed through skin.
- Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling.
- Avoid inhaling spray mist.
- Do not use or store near food or feed.

See side panel for additional precautionary statements and statement of practical treatment.

STORE IN A COOL, DRY PLACE.
 STIR THOROUGHLY TO MIX CONTENTS BEFORE USING.
 5 GAL. NET

Made in U.S.A. by
STAUFFER CHEMICAL COMPANY
 WESTPORT, CT 06880

SEED TREATMENT DOSAGE RATES

SEED	DOSAGE RATES
Alfalfa, clover, lespedeza, trefoil	5.8 1/4 fl. oz /100 lb.
Beans (snap, dry, cowpeas)	2.3 fl. oz /100 lb.
Beets (garden)	8 1/2 fl. oz /100 lb.
Cereal grains (for seed and seedling rots — not for bunt smut)	
Barley, rye	2.3 fl. oz /100 lb.
Oats	2.4 fl. oz /100 lb.
Wheat	11.4 fl. oz /100 lb.
Cole crops (broccoli, Brussels sprouts, cabbage, cauliflower)	1.2 fl. oz /100 lb.
Corn	
Field	11.4 2 1/4 fl. oz /100 lb.
Sweet	2.4 fl. oz /100 lb.
Cotton — Acid delinted	2.4 fl. oz /100 lb.
Reginned, fuzzy	2 1/2 fl. oz /100 lb.
Cucumber, cantaloupe	3 1/2 fl. oz /100 lb.
Crucifers (Mustard, radish, rape, turnips)	2 1/2 fl. oz /100 lb.
Flax	1.2 fl. oz /100 lb.
Grasses	2.3 1/4 fl. oz /100 lb.
Milo	5.8 1/4 fl. oz /100 lb.
Peanuts (unshelled)	4.6 fl. oz /100 lb.
Peas, Lentils	6 fl. oz /100 lb.
Peppers	2 1/2 fl. oz /100 lb.
Rice	2.3 fl. oz /100 lb.
Sorghum (hulled) For Kernel smut	3.6 fl. oz /100 lb.
Soybeans	11.4 1/2 fl. oz /100 lb.
Spinach	11.2 fl. oz /100 lb.
Squash, pumpkin, watermelon, muskmelon	4.6 fl. oz /100 lb.
Sugar beets — West	11.2 fl. oz /100 lb.
East	6 fl. oz /100 lb.
Sunflowers	12 fl. oz /100 lb.
Swiss chard	2.4 fl. oz /100 lb.
	10-12 fl. oz /100 lb.

Further specific information on the subject of seed treatment may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

STORAGE AND DISPOSAL:

- 1. PROHIBITIONS:** — Store in a cool, dry place. — Do not contaminate water, food, or feed by storage or disposal. — Open dumping is prohibited. — Do not reuse empty container.
- 2. PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticide or buried in a safe place away from water supplies.
- 3. CONTAINER DISPOSAL:** Rinse container and dispose in a landfill approved for pesticide containers, or bury in a safe place away from water supplies.
- 4. GENERAL:** Consult Federal, State or local disposal authorities for approved alternative procedures.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE: Stauffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION: Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties; THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

EPA Reg. No. 476-2192
 EPA Est. No. 476-NB-1

BEST DOCUMENT AVAILABLE