(Stautter Shield)

1.

40 *

7³, **4**

一日本のないないである。そのためのないであるとうないであるとうないであるとうないであるというであるというないであるというないであるというないであるというないであるというないであるというないであるというない

等题

Country

Hex

United

ý.

States

 $I M I D A N \Theta I - E$

HOME GARDEN INSECTICIDE An Emulsifiable Concentrate

For control of insects specified in the directions on deciduous shade trees, evergreens, ornamental flowering trees and shrubs and roses.

ACTIVE INGREDIENTS: Phosmet: N-(Mercaptomethyl)phthalimide S-(0,0-dimethyl . 11.7% ACCIPTED 82.9% Petroleum Oil INERT_INGREDIENTS:.... 5.4% • 10 U.S. 5 T00.0% CONTAINS 1 POUND IMIDAN PER GALLON FEP D KEEP OUT OF REACH OF CHILDREN Ur ' e e' Ti. WARNING STATEMENT OF PRACTICAL TREATMENT _ 476-2217 all a Physician Immediately: If a known exposure occurs or if poisoning is Spected, do not wait for symptoms to develop. Immediately initiate the recommended procedures described on the left panel of this label. Simultaneously contact a physician, the nearest hospital or a Poison Control Center, or call collect, day or night: (800) 424-9300 (203) 226-6602 OR STAUFFER CHEMICAL COMPANY PESTICIDE TEAM SAFETY NETWORK Inform the person contacted of the type and extent of exposure, describe the victim's symptoms and follow the advice given. NOTE: Be sure to advise the physician that the compound is a cholinesterase inhibitor and follow the physician's advice. NOTE TO PHYSICIAN: Exposure may cause cholinesterase inhibition. Atropine by injection is antidotal. 2-PAM is also antidotal when administered early and in conjunction with atropine. IF_SWALLOWED: Do NOT induce vomiting. Immediately give large quantities of wher and get medical attention. If vomiting occurs, give fluids again. Never give anything by mouth to an unconscious person. IF INHALED: Remove victim to fresh air. If not breathing, give artifical respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen, preferably with a physician's advice. IF ON SKIN: Flush with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Get medical attention if irritation occurs. Wash clothing before re-use. IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Call a physician. SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS •••• EPA Reg. No. 476-EPA Est. 476-GAL. NET RS-821028 This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified:

> Made in U.S.A. by STAUFFER CHEMICAL COMPANY WESTFORT, CT. 06881-0850

144

176-277

2.

IMIDAN® 1-E HOME GARDEN INSECTICIDE

(LEFT PANEL)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

WARNING:

- Causes eye irritation.
- May cause skin irritation.
- May be harmful if swallowed or inhaled. Avoid breathing spray mist.
- Do not get in eyes, on skin or on clothing.
- In case of contact, flush eyes with plenty of water. For skin, wash with soap and water. Wash contaminated clothing with soap and hot water before re-use. Wash thoroughly after handling.
- Do not contaminate food or feed.

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.
- This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

STORAGE AND DISPOSAL

- KEEP OUT REACH OF CHILDREN.
- Protect from temperatures below 25°F. Product crytallizes at lower temperatures. Warm or store at higher temperatures, and mix to redisolve crystals and assure uniformity before use.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container

		4
IMIDAN® 1-E HOME GARDEN INSECTIO	CIDE	3.
	(RIGHT PANEL)	
	PRODUCT INFORMATION	
• MIX WELL BEFORE U	SING.	
 IMIDAN 1-E is an applications. 	emulsifiable liquid for u	se in water for spray
 Read all label di only as specified 	rections before using. A on this label.	pply this product
 Keep children and 	pets away from immediate	area while spraying.
• Do not apply duri	ng periods of abnormally	high humidity.
• Do not overdose.		
INSECTS CONTROLLED	BY IMIDAN 1-E ARE THE FOL	LOWING:
Birch Leafminer Elm Leaf Beetle Elm Spanworm European Red Mite Fall Cankerworm Fall Webworm Forest Tent Caterpi Fruit Tree Leaf Rol Green Apple Aphid Gypsy Noth Japanese Beetle <u>RECOMMENDATIONS</u> : IMIDAN 1-E may be u <u>Shade Trees</u> Ash Beech Birch	Leaf Tier Maple Aph Mealy Bug Mimosa We Pine Bark Rose Aphi Spring Ca Two-Spott Willow Le Woolly Ap Woolly Be Sed on the following plan Elm Locust Maple	id bworm Aphid d nkerworm ed Mite af Beetle ple Aphid ech Aphid
Flowering Trees a Dogwood Euonymus Lilac Jasmine Magnolía	nd Shrubs	on Washington Hawthorne-
Evergreens Hemlock Holly Juniper Pine Rhododendron Azalea Pyracatha	Fir Spruce Arborvitae Mountain Laurel Cotoneaster Boxwood Yew	

to be starting as a second

ł,

まってい

Ν.

;

•____

- 2 ,

177

٦

_

.

÷

IMIDAN⊕ 1-E HOME GARDEN INSECTICIDE

4

4. 476-2217

- n' 🗯

1.

*****۵.

DIRECTIONS FOR USE

It is ~ violation of Federal law to use this product in a manner inconsistent with its labeling.

• Mix well before using.

- . As a general foliar spray on shade trees and ornamentals, use 2 to 6 teaspoonsful per gallon in a pump-up sprayer or small estate sprayer or a hose-end sprayer. Apply thoroughly to all plant surfaces when insects appear. Repeat as necessary.
- Pour the recommended amount of IMIDAN 1-E into the spray tank as you add the water.
- Improved wettability can be achieved with the addition of a spreader/sticker.

DOSAGE RATES

INSECTS		<u>_S</u>	PRAYER DILUTIONS
Green Apple Aphid Maple Aphid Rose Aphid Woolly Apple Aphid Woolly Beech Aphid	Use	2	to 4 teaspoonsful per gallon.
Elm Leaf Beetle Elm Spanworm Fall Cankerworm Fall Webworm Forest Tent Caterpillar Gypsy Moth Japanese Beetle Spring Cankerworm Willow Leaf Beetle			to 6 teaspoonsfull per gallon. teaspoonsful for early instars.
Birch Leafminer European Red Mite Mealy Bug Pine Bark Aphid			teaspoonsful per gallon.
Fruit Tree Leafroller Mimosa Webworm Two-Spotted Mite	Use	6	
NOTICE: Stauffer Chemical implied, including the warr	Comp rante purp	an es os	<pre>y makes no warranty, express or * of merchantability and/or e, concerning this material,</pre>

(Stautter Shield)

IMIDAN_O1-E

HOME GARDEN INSECTICIDE An Emulsifiable Concentrate

For control of insects specified in the directions on deciduous shade trees, evergreens, ornamental flowering trees and shrubs and roses. ACTIVE INGREDIENTS:

Phosmet: N-(Mercaptomethyl)phthalimide S-(0,0-dimethyl INERT INGREDIENTS:.... 5.4% • • • • • • 100.0% CONTAINS 1 POUND IMIDAN PER GALLON KEEP OUT OF REACH OF CHILDREN Unit - Co F. WARNING represed to a STATEMENT OF PRACTICAL TREATMENT 476-2217 (all a Physician Immediately: If a known exposure occurs or if poisoning is spected, do not wait for symptoms to develop. Immediately initiate the recommended procedures described on the left panel of this label. Simultaneously contact a physician, the nearest hospital or a Poison Control Center, or call collect, day or night: (203) 226-6602 (800) 424-9300 STAUFFER CHEMICAL COMPANY PESTICIDE TÉAM SAFETY NETWORK Inform the person contacted of the type and extent of exposure, describe the victim's symptoms and follow the advice given. NOTE: Be sure to advise the physician that the compound is a cholinesterase inhibitor and follow the physician's advice. NOTE TO PHYSICIAN: Exposure may cause cholinesterase inhibition. Atropine by injection is antidotal. 2-PAM is also antidotal when administered early and in conjunction with atropine. IF SWALLOWED: Do NOT induce vomiting. Immediately give large quantities of w er and get medical attention. If vomiting occurs, give fluids again. Never give anything by mouth to an unconscious person. IF INHALED: Remove victim to fresh air. If not breathing, give artifical respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen, preferably with a physician's advice. IF ON SKIN: Flush with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Get medical attention if irritation occurs. Wash clothing before re-use. IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Call a physician.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 476-EPA Est. 476____GAL. NET__L RS-821028

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States

Made in U.S.A. by STAUFFER CHEMICAL COMPANY WESTPORT, CT. 06881-0850 144

196 1911

IMIDAN⊕ 1-E HOME GARDEN INSECTICIDE

(LEFT PANEL)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

WARNING:

- Causes eye irritation.
- May cause skin irritation.
- May be harmful if swallowed or inhaled. Avoid breathing spray mist.
- Do not get in eyes, on skin or on clothing.
- In case of contact, flush eyes with plenty of water. For skin, wash with soap and water. Wash contaminated clothing with soap and hot water before re-use. Wash thoroughly after handling.
- Do not contaminate food or feed.

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.
- This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

STORAGE AND DISPOSAL

- KEEP OUT REACH OF CHILDREN.
- Protect from temperatures below 25°F. Product crytallizes at lower temperatures. Warm or store at higher temperatures, and mix to redisolve crystals and assure uniformity before use.

• Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry place. Do not store near sources

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

.....

112. -- 10

3.

IMIDAN® 1-E HOME GARDEN INSECTICIDE

(RIGHT PANEL)

PRODUCT INFORMATION

. MIX WELL BEFORE USING.

- IMIDAN 1-E is an emulsifiable liquid for use in water for spray applications.
- Read all label directions before using. Apply this product only as specified on this label.
- Keep children and pets away from immediate area while spraying.

Do not apply during periods of abnormally high humidity.

Do not overdose.

INSECTS CONTROLLED BY IMIDAN 1-E ARE THE FOLLOWING:

Birch Leafminer Elm Leaf Beetle Elm Spanworm European Red Mite Fall Cankerworm Fall Webworm Forest Tent Caterpillar Fruit Tree Leaf Roller Green Apple Aphid Gypsy Moth Japanese Beetle Leaf Tiers Maple Aphid Mealy Bug Mimosa Webworm Pine Bark Aphid Rose Aphid Spring Cankerworm Two-Spotted Mite Willow Leaf Beetle Woolly Apple Aphid Woolly Beech Aphid

RECOMMENDATIONS:

IMIDAN 1-E may be used on the following plant species:

Shade Trees	
Ash	Elm
Beech	Locust
Birch	Maple

Oak Willow Hickory

Flowering Trees and Shrubs Dogwood Hawthorne (Do not use on Washington Hawthorne-Euonymus injury may result). Lilac Crepe Myrtle Jasmine Forsythia

MagnoliaPriveEvergreens
HemlockFHollySJuniperAPineMRhododendronCAzaleaBPyracathaY

Hickory

. •

4

. . .

•

••••

••••

rivet			
Fir			
Spru	ce		
	rvita	1 e	
Moun	tain	Laurel	
Coto	neast	ter	
Boxw	ood		
Yew			

Roses

INIDAN⇔ 1-E HOME GARDEN INSECTICIDE

476-)217 4.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

• Mix well before using.

- As a general foliar spray on shade trees and ornamentals, use 2 to 6 teaspoonsful per gallon in a pump-up sprayer or small estate sprayer or a hose-end sprayer. Apply thoroughly to all plant surfaces when insects appear. Repeat as necessary.
- Pour the recommended amount of IMIDAN 1-E into the spray tank as you add the water.
- Improved wettability can be achieved with the addition of a spreader/sticker.

DOSAGE RATES

INSECTS		<u>_</u> <u>S</u>	PRAYER_DILUTIONS
Green Apple Aphid Maple Aphid Rose Aphid Woolly Apple Aphid Woolly Beech Aphid			to 4 teaspoonsful per gallon.
Elm Leaf Beetle Elm Spanworm Fall Cankerworm Fall Webworm Forest Tent Caterpillar Gypsy Moth Japanese Beetle Spring Cankerworm Willow Leaf Beetle	Use Use	4	to 6 teaspoonsfull per gallon. teaspoonsful for early instars.
Birch Leafminer European Red Mite Mealy Bug Pine Bark Aphid	Use	4	teaspoonsful per gallon.
Fruit Tree Leafroller Mimosa Webworm Two-Spotted Mite	Use	6	teaspoonsful per gallon.
NOTICE: Stauffer Chemical implied, including the warr	Comp rante purp	any es os(y makes no warranty, express or of merchantability and/or e, concerning this material,

EPA REGISTRATION NO. DATE OF ISSUANCE U.S. ENVIRONMENTAL PROTECTION AGENCY 0.1 OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (#11-567) TERM OF ISSUANCE WASHINGTON, D.C. 20460 NAME OF PESTICIDE PRODUCT NOTICE OF PESTICIDE: (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended) NAME AND ADDRESS OF REGISTRANT (Include ZIP code) 285 1/11/83 1/11/83 Γ Т BEST DOCUMENT AVAILABLE L NGTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. * the tabeling accepted in connection with this Registration/Reregistration is returned herewith. A cr is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect 1. 1940 a the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pest-icid an accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. 1-ATTACHMENT IS APPLICABLE SIGNATURE OF APPROVING OFFICIAL DATE

EPA FOR 8570-6 (Rev. 5-76) PREVIOUS FOILION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.