CORRECTED 10/23/69

FARMRITE DRY POWDERED ARSENATE OF LEAD

ACTIVE INGREDIENTS:

Dry Lead Arsenate - - -_ _ _ _ _ _ _ _ _ _ _ _ _ _ _ INERT INGREDIENTS: - - - - -2.01

Total - - - - - - - 100.0 (

Total Arsenic: Expressed as Metallic (AS) - - - - - 20.5% Total Lead: Expressed as metallic (Pb) - - - - - 58.0% Arsenic in water soluble form: Expressed as Metallic (AS) 0.35%

> X (printed in red on contrasting O POISON C background)

> > DANGER (18 point type)

KEEP OUT OF REACH OF CHILDREN (12 point type)

Poisonous if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid breathing dust or spray mist.

ANTIDOTE: Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Then give 2 tablespoonfuls of Epsom Salts or Milk of Magnesia in water, and plenty of milk and water. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY.

DIRECTIONS


Add the required amount of lead arsenate to the partly filled spray tank with agitator running and fill tank to desired volume with water. If preferred, the lead arsenate may be mixed into a slurry with water in a bucket and then poured into the spray tank. Make complete coverage of fruit and foliage for best results.

RECOMMENDATIONS

APPLES: For control of Codling Moth, Plum Currulio, Red Banded Leaf Roller; use 2 to 3 pounds of lead arsenate in 100 gallons of spray along with other chemicals for insect and disease control. Begin lead arsenate application in petal fall spray and continue through fourth or fifth cover spray. Do not apply within 30 days of harvest. Remove excess residues at time of harvest. Do not graze treated areas or feed treated forage to livestock.

CHERRIES: For control of Cherry Maggot; use 2 to 3 pounds per 100 gallons of spray. Several cover sprays may be used if necessary. If used alone, add an equal amount of spray lime, as a safening material, to the spray tank. Do not apply within 30 days of harvest. Remove residues at time of harvest. Do not graze treated areas or feed treated forage to livestock.

TOBACCO: For control of Hornworms, use 4 pounds in 50 gallons of water and apply 50 to 30 gallons per acre. Begin applications when insects first appear and repeat as necessary.



***/** ·

.

Pane ? Cerrected 10/23/64

STADE TREES AND CREATERN LANT : Cluerpilling use d to b coundry the top of a line of the second sec • norsec by the complete me should be

FARMPITE DIY POWDERED ARGENATE OF THAT


M nufactured in-

CENTRAL CHEMICAL CORPORATION

General Offices Hagesstown, Maryland

____ 1bs. Net

USDA Reg. no. 477-18



6 6 1 1 6 1 1 8 9 1 1 8 9 1