

# Farnatho Copper

A blend of tribasic copper sulfate and neutral carriers to be used as a fungicide dust on fruits and vegetables.

**ACTIVE INGREDIENTS:**

Copper (as metallic) 1.00%

**INERT INGREDIENTS:** 99.00%

Total 100.00%

### CAUTION

Harmful if swallowed.

### KEEP OUT OF REACH OF CHILDREN

Burn bag immediately when empty. Stay out of smoke.

### RECOMMENDATIONS


**POTATOES AND TOMATOES:** Early and late blight.—Use 30 to 40 pounds per acre. Begin applications when plants are 4 to 8 inches high and repeat at 7 to 10 day intervals. Dust thoroughly the under and upper sides of leaves.

**CELERY:** BLIGHT, SEPTORIA LEAF SPOT.—Use 30 to 40 pounds per acre. Begin applications when disease first appears and repeat at 7 to 10 day intervals until harvest.

**CARROTS:** Early and late blight.—Use 30 to 40 pounds per acre. Begin applications when disease first appears and repeat at 7 to 10 day intervals until harvest.

**CUCUMBERS AND MELONS:** Downy mildew, anthracnose, leaf blight.—Use 30 to 40 pounds per acre. Begin applications when first leaves are fully developed. Dust thoroughly the under and upper sides of the leaves. Repeat at 7 to 10 day intervals.

**GRAPES:** Black rot, downy mildew.—Use 20 to 25 pounds per acre. Begin applications when leaves first appear. Repeat before blossoms open and after petal fall continuing at two week intervals until within two weeks of harvest.



Manufactured by

**CENTRAL CHEMICAL CORPORATION**

General Offices

Hagerstown, Maryland

NET WT. 50 LB.

FARMRITE MICROSUL WETTABLE SULFUR  
Page 2

Manufactured for  
CENTRAL CHEMICAL CORPORATION

General Offices  
Hagerstown, Maryland

NET WEIGHT \_\_\_ LBS.

U.S.D.A. reg. no. 477-57