



Methoxychlor 2-E

DIRECTIONS

Pour the recommended amount of this material into the partly filled spray tank. Add the remainder of the water to fill the tank. Keep agitator running during filling and spraying operations. Do not let the mixture stand. If the concentrate does not mix readily with the water but tends to separate as an oily layer, do not use as injury to plants may result.

RECOMMENDATIONS

FORAGE CROPS: Alfalfa, clover, grass, soybeans, peanut forage — Use the following amounts in sufficient water for thorough coverage of one acre and repeat at 7 to 14 day intervals as necessary. — **Leafhoppers, spittle bugs;** use 1 to 2 quarts. **Alfalfa caterpillars, flea beetles;** use 1½ to 2 quarts. **Alfalfa weevil larvae, Alfalfa webworm, fall armyworm, Clover leaf weevil, Mexican bean beetle, Pea weevil, Soybean caterpillar, Velvet bean caterpillar;** use 2 to 3 quarts. **Coupea Curculia, Japanese beetle;** use 3 quarts. **Army worm;** use 3 to 4 quarts. Do not apply within 7 days of harvest. As an alternate program for Alfalfa weevil larvae and as a control for pea aphid, use 2 to 3 quarts Methoxychlor 2-E plus 1 to 1½ pints of Malathion 5-E. Do not apply within 7 days of harvest.

POTATOES: Colorado potato beetle, leafhoppers, flea beetles — use 2 quarts in sufficient water for thorough coverage of one acre. Repeat as necessary.

LIVESTOCK: (except dairy animals) For control of certain external parasites. USE AS A SPRAY ONLY. DO NOT USE DIPPING VATS.

CATTLE: For control of shortnosed and long-nosed cattle lice, apply as a drenching spray; for control of horn flies, apply 2 quarts of spray per mature animal. To make spray mixture, use 1 quart Methoxychlor 2-E to 12½ gallons of water.

HOGS: For control of lice, use 1 quart Methoxychlor 2-E to 12½ gallons of water applied as a drenching spray.

An emulsifiable liquid for use in water as an insecticide. Contains 2 pounds Technical Methoxychlor per gallon.

ACTIVE INGREDIENTS:

Methoxychlor, Technical* 24.0%
Xylene Range Aromatic Hydrocarbon Solvent 69.0%

INERT INGREDIENTS

Total 100.00%

*24% Methoxychlor Technical is equivalent to 21% 2,2-bis (P-Methoxyphenyl)-1,1,1-trichloroethane and 3% other isomers and reaction products.

CAUTION

Avoid contact with skin. Avoid breathing dust or spray mist. Avoid contamination of feed and foodstuffs. This product is poisonous to fish. Keep out of any body of water. Fish may be killed.

KEEP OUT OF REACH OF CHILDREN

FLAMMABLE MIXTURE. KEEP AWAY FROM HEAT AND OPEN FLAME. If exposed to prolonged cold, place in warm storage (50° F. to 80° F.) for several hours and agitate before using.

Manufactured by

CENTRAL CHEMICAL CORPORATION

General Offices

Hagerstown, Maryland

Net Contents Gallons

SELF APPLICATOR FLY CONTROL ON CATTLE: (except dairy animals): For the control of horn flies, dilute 1 gallon Methoxychlor 2 E with 4 gallons of diesel fuel or furnace type oil. Do not use any other type of oil. Use 1 gallon of solution to saturate burlap on "back rubber", soak burlap every two to three weeks.

LARGE TYPE GRAIN STORAGE BINS: To clean up infestation of cadelle, flat grain beetle, saw toothed grain beetle, lesser grain borer, red flour beetle, long headed flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus beetle — use 1 gallon Methoxychlor 2-E in 10 gallons of water applied at the rate of 2 gallons of spray per 1,000 square feet before products are stored. NOTE: Do not add grain to bin for at least 24 hours or until walls have dried out thoroughly.

CONTACT AND SPACE SPRAY FOR FLIES: To reduce populations of house and stable flies in farm buildings, in barnyards, in alleys, on dumps, and other out door locations, dilute 1 pint Methoxychlor 2-E in 2½ gallons of water and apply in space by mist blower or directly to flies by hydraulic sprayer, hand sprayer or cylindrical pressure sprayer. Care should be taken to avoid contamination of milk handling equipment, feed stuffs, feed troughs, and water receptacles. Dairy animals should not be present while spraying.

NOTE: Do not use with any spray materials which are incompatible with oil. Do not apply within 14 days of sulfur or sulfur product applications.

WARRANTY

Seller makes no warranty of any kind, express or implied, concerning the use of this material. Buyer assumes all risk of use or handling whether in accordance with directions or not.

U.S.D.A. reg no. 477-152