

REG. OF. AND P. IN THE ACT  
 FEDERAL POISON REGISTER  
 SUBJECT  
 COMMENTS

# FARMRITE 5% SEVIN DUST

An insecticide dust containing Sevin\*

\*Trademark of Union Carbide Corporation for Carbaryl (l-naphthyl N-methylcarbamate)

**ACTIVE INGREDIENT:**

Carbaryl (l-naphthyl N-methylcarbamate) .....	5.0%
INERT INGREDIENTS: .....	95.0%
<hr/>	
Total .....	100.0%

## CAUTION

Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid storage near feed or food products.

### KEEP OUT OF REACH OF CHILDREN

#### DIRECTIONS FOR USE

**BEANS:** For Mexican bean beetle—apply Farmrite 5% Sevin Dust at the rate of 10 pounds per acre. For leafhoppers—apply 20 pounds per acre. For bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle and velvet bean caterpillar—apply 20 pounds per acre. For corn earworm, armyworms, stink bugs, tarnished plant hoppers—apply 20 to 40 pounds per acre. May be applied up to and including the day of harvest. When label directions are followed, treated crops may be fed to meat and dairy animals.

**CATTLE OR FIELD CORN:** For corn earworm—apply 10 to 40 pounds per acre, depending upon field conditions. Repeat at 2 to 5 day intervals throughout growing season. Direct dust on plants to and over weeds, etc., are thoroughly covered. Do not apply until 10 days before the day of harvest. Allow 10 days before harvest or marketing of produce.

**TOMATOES, EGGPLANT, PEPPER:** For tomato fruit worm, fall armyworm, flea beetle and leafhopper—apply 20 to 40 pounds per acre. Tomato hornworm, European cornborer and stink bug—apply 20 to 80 pounds per acre. May be applied up to 10 days before harvest.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI:** For flea beetle and harlequin bug—apply 10 to 20 pounds per acre. For corn earworm, armyworm, cabbage caterpillars—apply 20 to 40 pounds per acre. When cabbage looper is the principal problem, use an alternative pesticide. (For small loopers, after heads form, use Sevin every 5 to 7 days.) Apply at 7 to 10 day intervals. Do not apply within 3 days of harvest. When label directions are followed, treated crop residue may be fed to dairy and meat animals.

**POULTRY (chickens, turkeys, ducks, geese, game birds and pigeons):** HAND APPLICATION TO BIRDS: For control of Northern fowl mite, chicken mite, lice and fleas—use 1 pound to treat 100 birds. Apply with shaker can, squeeze bottle, or hand duster. Direct dust to nest and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and flea is a supplement to litter treatment for control of these pests. Do not treat birds within 7 days of slaughter.

per acre. Sevin may also be used to control DDT-resistant leafhoppers. May be applied up to and including the day of harvest.

**POTATOES:** For Colorado potato beetle, leafhopper and flea beetle—apply 10 to 20 pounds per acre. For egg bugs—apply 20 to 40 pounds per acre. Apply when insects or their damage first appear. Repeat at 7 to 10 day intervals or as necessary. May be applied up to and including the day of harvest.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI:** For flea beetle and harlequin bug—apply 10 to 20 pounds per acre. For corn earworm, armyworm, cabbage caterpillars—apply 20 to 40 pounds per acre. When cabbage looper is the principal problem, use an alternative pesticide. (For small loopers, after heads form, use Sevin every 5 to 7 days.) Apply at 7 to 10 day intervals. Do not apply within 3 days of harvest. When label directions are followed, treated crop residue may be fed to dairy and meat animals.

**POULTRY (chickens, turkeys, ducks, geese, game birds and pigeons):** HAND APPLICATION TO BIRDS: For control of Northern fowl mite, chicken mite, lice and fleas—use 1 pound to treat 100 birds. Apply with shaker can, squeeze bottle, or hand duster. Direct dust to nest and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and flea is a supplement to litter treatment for control of these pests. Do not treat birds within 7 days of slaughter.

**LITTER AND ROOST TREATMENT:** Apply 1 pound per 40 square feet to control chicken mite, fleas and bed bugs. Use scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and water troughs.

Caution: Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter.

**TOBACCO— IN PLANT BEDS:** For tobacco flea beetle—apply 1 to 2 pounds per 100 square yards, when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplanting. **IN FIELDS:** For hornworm and tobacco flea beetle—apply 15 to 25 pounds per acre. Good coverage is essential. Repeat as needed. Use highest dosage on large plants. Avoid heavy applications just before pruning leaves. If late applications are necessary, treat immediately after pruning rather than before pruning. Treated plants may be entered immediately after dust has settled.

**ORNAMENTALS:** Arbor vitae, aster, azalea, birch, boxwood, camellia, carnation, chrysanthemum, dahlia, dogwood, elm, gladiolus, holly, juniper, lilac, marigold, petunia, rhododendrum, rose, rose of Sharon, yew, zinnia. For control of certain aphids, bagworm, blister beetle, boxelder bug, certain leaf miners, elm leaf beetle, flea beetle, Japanese beetle, leafhopper, leaf rollers, orange tortrix, periodical cicada, plant bug, scale insects, tent caterpillar, spotted thrips—dust thoroughly to cover the upper and under sides of the foliage with Farmrite 5% Sevin Dust.

#### WARRANTY

Under the terms of a written warranty, we warrant, concerning the use of this product, that it will control the insects and diseases mentioned on the label when directions are followed.

Manufactured by

## CENTRAL CHEMICAL CORPORATION

General Office

Hagerstown, Maryland

**Net Weight 50 Lbs.**

Form No. A-1 (Rev. 1-1-55)