

2/16/66

FARWRITE 15% ZINEB DUST

ACTIVE INGREDIENTS:

Zineb (zinc ethylene bis(ithi) carbonate) ----- 15.0%
Metallic Zinc Content 4.1%

INERT INGREDIENT : ----- 85.0%

Total 100.0%

CAUTION (10 - int type)

May cause irritation of nose and throat. Do not breathe dust or spray mist.

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS

Farwrite 15% Zineb Dust may be applied from this package with a suitable duster at any time during the day to control the following diseases of mushrooms: verticillium disease (brown spot, dry bubble) dactylium (mildew, soft decay, cobweb) and mycogone disease (bubbles).

Farwrite 15% Zineb Dust may be used at the rate of 0.5 pounds and 0.7 pounds per 4,000 square feet. The first dust application should be made soon after the casing soil is applied and continue applications at 3 to 7 day intervals until small buttons begin to form. Continue applications between breaks as long as needed. Care should be taken not to apply this dust directly on the mushrooms. Consult your county agent for additional information regarding the use of this dust on other crops.

WARRANTY

Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risk of use and handling whether in accordance with directions or not.

Manufactured by

CENTRAL CHEMICAL CORPORATION

General Offices
Hagerstown, Maryland

NET 50 LBS.

U.S. Reg. No. 477-202

RECEIVED
477-202