

to 18 inches, apply 1 gallon of mixture per 5 lineal feet. Outside of foundation walls, including entrance platforms and porches, apply 2 gallons of mixture per 5 lineal feet of trench where the foundation is not more than 15 to 18 inches deep. If more than 15 to 18 inches deep but not over 30 inches deep, apply 4 gallons per 5 lineal feet. As a pre-construction treatment, apply 1 gallon per 10 square feet of soil surface where attached porches, entrance platforms, utility entrances, etc., have covering slab on fill ground. A minimum of 6 weeks control is obtained by use of a 1% Chlordane emulsion in above applications. NOTICE: Do not use on dairy animals or animals being fattened for slaughter. Avoid drift onto adjoining food crops and forage crops. To protect fish and wild life, do not contaminate lakes, ponds, or streams. Use only as recommended on the label.

WARRANTY

Under no circumstances is there any warranty of any kind, express or implied, concerning the use of this material. Buyer assumes all risk of use or handling whether in accordance with directions or not.

Net Contents Gallons

One gallon per 10 sq. ft.

U.S.D.A. reg. no. 477-210

Lawn and Turf

ANTS: Use 1 pint in 50 gallons of water and drench the colony. CHIGGERS AND TICKS: use 1 to 1½ pints in sufficient water to cover an acre. Asiatic Garden Beetle larvae, European chafer larvae, Green June beetle larvae, Japanese beetle larvae, White fringed beetle larvae, White grubs, Wireworms use 2½ pints to 5 pints in sufficient water for uniform coverage of one acre. Wash into top few inches of soil. Do not allow children or pets on treated area until Chlordane has been watered down and the treated area has dried throughly. Birds feeding on treated areas may be harmed.

TERMITE CONTROL

IN BUILDINGS Slab type construction: As a pre-construction treatment under the slab and under attached porches, use ½ gallon in 18 gallons of water and apply 1 gallon of mixture per 10 square feet as an overall treatment. For existing houses, apply 2 gallons of above mixture per 5 lineal feet along the outside of the foundation using a trench to the depth of the foundations. For crawl space houses and houses with basements, apply 2 gallons of the above mixture per 5 lineal feet as a trenching application to critical areas under the house along the inside of foundation walls and around piers and utility entrances. If trench is deeper than 15



FARMER'S

CHLORDANE

An emulsifiable concentrate for
ACTIVE INGREDIENTS:

Technical Chlordane*

Petroleum Solvent

INERT INGREDIENTS:

* Equivalent to 44.4% Octachlorocyclopentadiene
29.6% related compounds
Contains 8 pounds Chlordane per gallon

CALIFORNIA

Harmful if swallowed. Contact with breathing spray mist. In case of contact with eyes, flush with water. Avoid contamination of feed and food. Wildlife in contact with treated area may be harmed if their waters are contaminated with Chlordane.

KEEP OUT OF REACH OF CHILDREN

Manufactured by


CENTRAL CHEMICAL

General Offices

477-210
ACCEPTED
Mar 14 1967
FEDERAL
EPA
PUNISH
UNDER NO
ATTACHED COMMENTS



FARMRITE



CHLORDANE 8-E

An emulsifiable concentrate for use in water as an insecticide.

ACTIVE INGREDIENTS:

Technical Chlordane*	74.0%
Petroleum Solvent	21.0%
INERT INGREDIENTS:	5.0%

Total 100.0%

* Equivalent to 44.4% Octachloro-1,7 methanotetrahydroindane and 29.6% related compounds.

Contains 8 pounds Chlordane per gallon.

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Wildlife in contact with treated areas may be harmed. Fish may be killed if their waters are contaminated with this product.

KEEP OUT OF REACH OF CHILDREN

Manufactured by

CENTRAL CHEMICAL CORPORATION

General Offices

HAGERSTOWN, MARYLAND

Lawn and Turf

1 pint in 50 gallons of water and trench
BUGGERS AND TICKS: use 1 to 1½
water to cover an acre. Asiatic Gard
European chafer larvae, Green June
panese beetle larvae, White fringed beetle
ubs, Wireworms use 2½ pints to 5
nt water for uniform coverage of one
p top few inches of soil. Do not allow
on treated area until Chlordane has been
nd the treated area has dried thoroughly.
on treated areas may be harmed.

TERMITE CONTROL

Slab type construction. A pre
atment under the slab and under at
use ½ gallon in 48 gallons of water and
of mixture per 10 square feet as an
t. For existing houses, apply 2 gallons
e per 5 linear feet along the outside of
sing a trench to the depth of the founda
wl space houses and houses with bas
gallons of the above mixture per 5 linear
hing application to critical areas under
the inside of foundation wall and around
r entrance. If trench is deeper than 1



ONIONS: Thrips use ¾ quart per 100 g
water per acre. Do not apply to green or sprin

TOMATOES: Flea beetle, Serpentine leaf m
mato hornworm use ½ to 1 quart to 100 g
water per acre. Repeat as necessary. Do not ap
fruit begins to form.

Soil Applications

WHITE GRUBS: Potatoes, strawberries, sugar
use 1½ to 2½ quarts per acre in 100 gallons of
soil surface as a pre-plant application. Disk
top 3 to 4 inches of soil immediately.

WIREWORMS: Beans, beets, cauliflower, onio
toes, sugar beets, sweet potatoes, turnips u
2½ quarts in 100 gallons of water per acre. Ap
surface as a pre-plant application. Disk into t
inches of soil immediately.

Shade Trees and Ornamentals

For Bagworm, Leaf beetles, Elm leaf beetles,
er, Lace bugs, Tree crickets on Arborvitae, Cam
Hawthorn, Holly, Junipers, Magnolia, Maple an
use 1 pint in 100 gallons of water. Begin ap
in late spring or when insects first appear.
2 to 3 weeks when necessary.

- PLUS - 100% CONTROL OF FLORAWEED, GROUND RY, KNOTWEED, HENBIT, PURSLANE, BLACK MEDIC, AND ALL OTHER COMMON WEEDS
- PLUS - 100% CONTROL OF ALL INSECTS PLUS THE 100% MAJOR AND TRADE ELEMENTS OF ALL INSECT DAMAGE
- PLUS - 100% CONTROL OF ALL WEEDS
- PLUS - 100% CONTROL OF ALL INSECTS, INCLUDING ALL OTHER INSECT DAMAGE AND ALL INSECTS

W

WEED and FEED PLUS

12 - 4 - 8

CAUTION

KEEP OUT OF REACH OF CHILDREN

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Sponholz (1980). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Sponholz (1980). The total carotenoid content was determined by the method of Arar and Cook (1980). The total carotenoid content was determined by the method of Arar and Cook (1980).

The authors are grateful to the anonymous referees and the editor for their valuable comments and suggestions. The authors also thank the participants at the 2012 Annual Meeting of the European Association of Agricultural Economists for their helpful comments. The authors are also grateful to the Department of Economics, University of Oxford, for providing the opportunity to work on this paper. The authors also thank the Department of Economics, University of Oxford, for providing the opportunity to work on this paper. The authors also thank the Department of Economics, University of Oxford, for providing the opportunity to work on this paper.

The present study of the hippocampal KLF10 promoter was the first to show that the promoter is active in the hippocampus and that the promoter is active in the hippocampus.

5. ABSTRACT, REFERENCE AND INSTRUCTION ANALYSIS

[illegible]

^a The number of subjects who were included in each group was 10.

NET WT. 22½ LBS.

PACKED BY

PACIFIC SUPPLY COOPERATIVE

PORTLAND, OREGON • SUNNYSIDE, WASHINGTON • HEYBURN, IDAHO

ACC

ds
\$3.95

WEED and FEED PLUS

PACIFIC WEED and FEED PLUS

ANALYSIS:

Total Nitrogen (N)	12.00
Available Phosphoric Acid P ₂ O ₅	4.00
Soluble Potash (K ₂ O)	3.00
Sulphur (combined S)	12.00
Zinc (ZN)	1.05
Iron (Fe)	2.00
Magnesium (Mg)	2.50
Calcium (Ca)	4.00


INGREDIENTS

DIRECTIONS:

Spread contents of this bag evenly over 3,000 sq. ft. (6 ft. x 6 ft. x 20 ft.).
Set spreader at low setting so that material will fall in a thin layer.
Run it to just pass through.
Spread on the bag is for use on lawns, golf courses, and other areas.
For more facts, ask your dealer or write to the manufacturer.

WEED

F



ACCEPTED

Soluble Potash (K ₂ O)	8.00%
Sulphur (combined S)	12.00%
Zinc (ZN)	.05%
Iron (Fe)	2.00%
Magnesium (Mg)	2.50%
Calcium (Ca)	4.00%

DIRECTIONS:

Spread contents of this bag evenly over 3,000 square feet of lawn. Set spreader at low setting so that the material will fall from the spreader like a pencil to just pass through.

Spread on the basis of 1/2 pound material per 100 square feet. This will cover 200 square feet. (A one pound coffee can level 100 square feet.) Spread on WEED and FEED-PLUS. See spreader settings below.

1. Do not apply to a new lawn, wait until lawn is three months old and has been mowed four times.
2. Do not mow lawn for two days prior to or two days after application. It is desirable to have maximum leaf surface for an effective application.
3. Apply to a DRY lawn, but preferably one that has previously been thoroughly moistened to a depth of two or three inches by rain or sprinkler. Weeds that have been hardened from growing in dry soil are not easily killed.
4. Take care to apply evenly with an accurate spreader and correct dial setting. Do not turn corners without shutting off spreader. If your spreader is not shown below, proceed with caution using a number smaller than half the largest number on the dial. For many spreaders with diamond, rectangular or oval holes an opening the size of an ordinary wooden pencil will give an approximate setting.
5. Do not water lawn for 24 hours after application. Should a heavy or prolonged rain occur, it may be necessary to re-apply for good weed control. Resume normal watering procedures after 24 hours.
6. Mouse ear chick weed and Japanese clover occasionally require two applications at the standard rate a month apart. Do not make double applications at one time.

SUGGESTED SPREADER SETTINGS

To apply 1/2 pound per 100 sq. ft. set spreader at	Lawn Beauty	Scott
	12 1/2	4 1/2

NOTE: Cyclone type spreaders should not be used. These settings were determined with new spreaders. Condition of your spreader, speed of walking and degree of fullness of spreader may vary output.

NOTICE

Pacific Supply Cooperative makes no warranty, express or implied, concerning this material, except that it conforms to the chemical description on the label. Neither Pacific Supply Cooperative nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on the condition