

11/15/67

FARMRITE 5.6% MANEB DUST

ACTIVE INGREDIENTS:

Maneb (Manganese ethylenebisdithiocarbamate) - - - - -	5.6%
(Manganese equivalent as metallic 1.00%)	
INERT INGREDIENTS: - - - - -	94.4%

Total - - - - - 100.0%

KEEP OUT OF REACH OF CHILDREN (12 point type)

DIRECTIONS FOR USE

Apply as a dust at 22½ to 37½ pounds per acre beginning when disease threatens being careful to treat all plant surfaces; repeat applications at 5 to 10 day intervals.

POTATOES: Early blight and late blight, use 30 to 37½ pounds of 5.6% Maneb Dust. Begin applications when plants are 2 to 6 inches high and continue at 5 to 10 day intervals as long as blight threatens.

TOMATOES: Early blight, late blight, spetoria leaf spot, gray leaf spot (stemphylium) and anthracnose - begin applications when fruits of first clusters are forming.


BEANS: Downy mildew - start applications of 22½ to 33-3/4 pounds per acre when disease first appears and repeat at 5 to 7 day intervals as long as disease threatens.

CELERY: Early blight and late blight - use 26-1/4 to 33-3/4 pounds of dust per acre beginning when plants are established in the field and continuing as frequently as required (every 3 to 5 days under severe blight conditions). Remove excess residues at harvest by stripping, trimming and washing.

CABBAGE: Alternaria leaf spot and downy mildew- use 21½ to 28½ pounds of dust per acre. Begin applications 7 to 10 days after planting. Repeat every 3 days until plants are set in the field or until field seeded plants are thinned. Do not apply within 7 days of harvest.

CUCUMBERS: Downy mildew and anthracnose - apply when disease threatens, repeat every 7 to 10 days.

MANUFACTURED BY



CENTRAL CHEMICAL CORPORATION

General Offices
Hagerstown, Maryland

NET _____ LBS.

USDA Reg. No. 477-X