

2/22/68

FARMITE POTATO SEED TREATER
(Contains Polycarb[®])

ACTIVE INGREDIENT:

Mixture of 0.2 parts by weight (84.9%) of ammoniated of
Zinc Polycarb (dithiocarbamate) zinc with 1 part by weight
(15.1%) ethylenebis /dithiocarbamic acid/ bimolecular and
trimolecular cyclic anhydrosulfides and aldisulfides - - - - - 7.1

INERT INGREDIENTS: - - - - - 92.9

100 %

Zinc expressed as metallic - 1.32%
U.S. Patent Pending

CAUTION (18 point type)

KEEP OUT OF REACH OF CHILDREN (12 point type)

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat
and skin! In case of contact, flush with plenty of water; for eyes, get medical
attention. Wear rubber gloves and respirator when mixing Seed Treater with potato
seed pieces.

DIRECTION FOR USE

Potato-Cut Seed: Fusarium Seed Piece Decay, Black Leg, Seed-borne Common Scab --
Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed pieces. For effective
control, Seed Treater must be applied to entire surface of each piece immediately
after cutting. If planting is delayed, follow the recommended storage practices of
cut seed pieces in your area.

Potato-Whole Seed: Seed-borne Common Scab - apply 1 to 1½ pounds of Seed Treater to
each 100 pounds of seed. Seed Treater should be applied evenly to entire surface.
If the treated whole seed is cut, a second application will be necessary if Black Leg
and Fusarium Seed Piece Decay are to be controlled.

[®] Polycarb is a registered Trademark of Badische Anilin & Soda Fabrik A.G.

MANUFACTURED BY

CENTRAL CHEMICAL CORPORATION

General Offices
Hagerstown, Maryland

NET _____ LBS.

REGISTERED
FEB 29 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REG-STER-
ED UNDER NO. 477-K SUBJECT
TO ATTACHED COMMENTS.

477-262

AC

USDA Reg. No. 477-K