



FARMRITE FOLPET 50-W



A Fungicide for Plant Disease Control

ACTIVE INGREDIENT:

(Folpet [N-(Trichloromethylthio)phthalimide])

INERT INGREDIENTS:

ACCEPTED
 NOV 5 1969
 477-275
 UNDER THE...
 FUNGICIDE...
 FOR ECON...
 ED UNDER...

50%
 50%
 100%



CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhalation of dust or spray mist. Avoid prolonged or repeated contact with skin. Avoid contamination of feed or foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product as specified on this label.

COMPATIBILITY

Folpet 50-W may be combined with most commonly used insecticides, adjuvants, fungicides. Fungicidal activity of Folpet is reduced if combined with strongly alkaline materials such as hydrated lime. This material should not be used in combination with or closely following oil sprays. Do not use with TEPP when a concentrate mixture is being applied.

DIRECTIONS FOR USE

Pour required amount of this material on surface of water in nearly filled spray tank, or pre-mix in a bucket half-filled with water and pour mixture through screen into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

State Agricultural Experiment Station or State Extension Service Specialist Consult for specific information as the timing, number and rate of applications needed will vary with local conditions.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

RECOMMENDATIONS

Unless otherwise stated, dosages are given in terms of pounds of Folpet 50-W per 100 gallons of water.

APPLES, CRABAPPLES: Bitter Rot, Botryospheria, Alternaria Spot, Brooks Fruit Spot, Sooty Blotch, Black Rot, Fly Speck and Scab - 1½ to 2 lbs. Apply up to 800 gallons per acre. Make first application at delayed-dormant and repeat at 7-14 day intervals until harvest. **CAUTION:** EAST-Use during the critical period (just before bloom until 30 days after petal fall) may cause russetting to sensitive varieties such as Golden Delicious, Red Delicious and Stayman Winesap.

AVOCADOS: Scab - 3 lbs. Make application at 2 to 3 week intervals as needed. **BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES:** Fruit Rot, Gray Mold, Spur Blight - 2 lbs. Apply 250 gallons per acre. Make first application when berries have set and repeat at 7 day intervals until harvest.

CANTALOUPE, MUSKMELONS, HONEYDEW MELONS, WATERMELONS, SUMMER SQUASH, PUMPKINS, WINTER SQUASH: Downy Mildew, Powdery Mildew, Anthracnose - 2 to 4 lbs. Apply up to 200 gallons per acre depending on size of vines. Make first application when first true leaves appear and repeat at weekly intervals until crop is harvested.

BLUEBERRIES, HUCKLEBERRIES: Gray Mold, Powdery Mildew, Leaf Spot Gloeosporium - 2 lbs. Apply 250 gallons per acre. Make first application before first infection. Repeat at 7 day intervals until harvest.

CELERY: Early Blight, Late Blight, Cercospora Leaf Spot - 2 lbs. Apply after plants are set in field and established. Repeat at 7 day intervals. Do not apply within 7 days of harvest.

CHERRIES (RED TART): Leaf Spot and Brown Rot - 1 to 2 lbs. At the 2 lb. per 100 gallon rate, do not use over 600 gallons per acre. Apply at pre-bloom and popcorn stage. Repeat in bloom and at 7 to 14 day intervals making last application to foliage post-harvest. Do not use on sweet cherry as foliage injury may occur.

CITRUS (ORANGES, GRAPEFRUIT, LEMONS, LIMES, TANGELOS, TANGERINES): Melanose, Scab, Greasy Spot - 2 lbs. Apply 300 gals., 2-3 petal fall, 2 weeks later and on the fall flush in August and September.

CRANBERRIES: Storage Rot - 9 lbs. per acre in sufficient water for thorough coverage. Apply during bloom and repeat at 10 to 14 day intervals. Do not apply within 30 days of harvest.

CUCUMBERS: Fruit Rots (Rhizoctonia and Pythium) - 6 to 8 lbs. per acre. Apply in a minimum of 200 gallons water as a directed spray at the soil in the row. Apply when plants are beginning to "run". Downy Mildew, Powdery Mildew, Anthracnose - 2 to 4 lbs. per acre. Add spray sticker according to manufacturer or State recommendations to the sprays.

GRAPES: Downy Mildew, Powdery Mildew, Black Rot, Dead Arm - 2 lbs. Apply 2 to 300 gallons per acre. Dead Arm - East - Make first application when new growth is 1 to 2 inches long, repeat when growth is 4 to 6 inches long. Black Rot, Downy Mildew - Make first application just before bloom, repeat immediately after bloom. Depending upon the susceptibility of the grape variety, from 1 to 3 more sprays may be needed at 7 to 10 day intervals. Powdery Mildew - East only - Make first application 2 to 3 weeks after bloom and repeat at 2-week intervals as needed. Note: During extended periods of hot dry weather, foliage injury to several varieties of grapes may occur.

GOOSEBERRIES, CURRANTS: Leaf Spot - 4 lbs. Apply 150 gallons per acre. Make first application just before bloom period. Repeat at 7 to 10-day intervals in the post-bloom period.

LETTUCE: Downy Mildew - 2 lbs. - Apply up to 200 gallons per acre. Make first application when plants are established in field. Repeat at 7-day intervals until harvest.

ONIONS, GARLIC, LEEKS, SHALLOTS: Purple Blotch and Downy Mildew - 4 lbs. Apply 200 gallons per acre. Make first application before first infection. Repeat at weekly intervals until tops begin to dry before harvest.

STRAWBERRIES: Leaf Spot, Fruit Rot - 2 lbs. Apply up to 200 gallons per acre. Make first application before bloom and repeat at 7 day intervals until fruit is harvested.

TOMATOES: Anthracnose - 4 lbs. Use up to 200 gallons per acre. Make first application during first bloom and repeat at 7 to 10-day intervals until fruit is harvested.

ROSES: Black Spot, Powdery Mildew, Rust - 2 lbs. per 100 gallons. (2 tablespoonfuls per 1 gallon of water). Make first application when young leaves first appear. Repeat at 7 to 10-day intervals or as necessary throughout the season. Make thorough application. Do Not Use Wetting Agents on Roses - Injury May Result.

CHRYSANTHEMUMS, IRIS: Sclerotinia Leafspot, Botrytis Leafspot - 2 lbs. per 100 gallons. (2 tablespoonfuls per 1 gallon of water). Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as needed throughout the season.

CARNATIONS: Rust, Alternaria Leafspot - 2 lbs. per 100 gallons (2 tablespoonfuls per 1 gallon of water). Begin application 2 weeks after planting and continue bi-weekly applications for 3 to 4 applications.

ZINNIAS, MARIGOLDS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS: Alternaria Leafspot, Powdery Mildew, Anthracnose - 2 lbs. per 100 gallons (2 tablespoonfuls per 1 gallon of water). Treat at least weekly during periods of rapid new growth and severe infection, and twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity.

POINSETTIA (GREENHOUSE): Pythium, Root Rot - 2 lbs. per 100 gallons water plus a spreader and use 1 pint per 8 inch pan. Make first application at time of panning. Repeat in 10 days.

AZALEA CUTTINGS: Damp-off - 1½ to 2 lbs. per 100 gallons. Allow to soak for 15 to 30 minutes before placing in rooting media. Destroy container by burying.

USDA Reg. no. 477-275

Manufactured For
CENTRAL CHEMICAL CORPORATION
General Offices Hagerstown, Maryland

..... LBS.