

477-279
ACCEPTED
Dec 23 1969
THE FEDERAL INSECTICIDE
ACT OF 1947

FARMRITE

COMP.

Farmrite Perthane[®] 4

EMULSIFIABLE CONCENTRATE

An agricultural insecticide containing 4 lbs. Technical Perthane per gallon.

ACTIVE INGREDIENTS:	
Diethyl diphenyl dichloroethane	46.6%
Related reaction products	2.4%
Xylene	47.0%
INERT INGREDIENTS:	
	4.0%
[®] Registered Trademark of Rohm & Haas	Total 100.0%



KEEP OUT OF REACH OF CHILDREN

CAUTION

Avoid contact with skin. Avoid contact with clothing. Wash hands after breathing spray mist. Avoid contamination of food and foodstuffs. Do not take internally. Harmful to fish and shellfish.

This product is toxic to fish. Keep out of lakes, streams, rivers. Apply only as directed on the label.

Empty containers should be kept out of reach of children. Do not reuse empty containers. Do not burn, melt, crush, or crush and burn in a safe place.

Directions for Use:

Measure the amount of this material and add to the sprayer tank. Mix thoroughly with the water before using. Stir contents thoroughly in hand sprayers. Do not use in areas where children, pets, or livestock are present. Do not use in areas where children, pets, or livestock are present.

APPLES: For control of apple maggot, apply 1.0 to 2.0 gallons of water per acre. For control of apple worm, apply 1.0 to 2.0 gallons of water per acre. For control of apple sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of apple borer, apply 1.0 to 2.0 gallons of water per acre. For control of apple moth, apply 1.0 to 2.0 gallons of water per acre. For control of apple fruitworm, apply 1.0 to 2.0 gallons of water per acre. For control of apple leafminer, apply 1.0 to 2.0 gallons of water per acre. For control of apple stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of apple stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

PEACHES: For control of peach twig borer, apply 1.0 to 2.0 gallons of water per acre. For control of peach fruitworm, apply 1.0 to 2.0 gallons of water per acre. For control of peach leafminer, apply 1.0 to 2.0 gallons of water per acre. For control of peach stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of peach stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

PLUMS: For control of plum curculionid, apply 1.0 to 2.0 gallons of water per acre. For control of plum fruitworm, apply 1.0 to 2.0 gallons of water per acre. For control of plum stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of plum stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

CHERRIES: For control of cherry fruit fly, apply 1.0 to 2.0 gallons of water per acre. For control of cherry stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of cherry stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

PEARL ONIONS: For control of pearl onion maggot, apply 1.0 to 2.0 gallons of water per acre. For control of pearl onion maggot, apply 1.0 to 2.0 gallons of water per acre.

SPINACH: For control of spinach leaf miner, apply 1.0 to 2.0 gallons of water per acre. For control of spinach leaf miner, apply 1.0 to 2.0 gallons of water per acre.

CRUCIFEROUS: For control of cabbage root fly, apply 1.0 to 2.0 gallons of water per acre. For control of cabbage root fly, apply 1.0 to 2.0 gallons of water per acre. For control of cabbage root fly, apply 1.0 to 2.0 gallons of water per acre.

PEACHES: For control of peach twig borer, apply 1.0 to 2.0 gallons of water per acre. For control of peach fruitworm, apply 1.0 to 2.0 gallons of water per acre. For control of peach leafminer, apply 1.0 to 2.0 gallons of water per acre. For control of peach stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of peach stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

PLUMS: For control of plum curculionid, apply 1.0 to 2.0 gallons of water per acre. For control of plum fruitworm, apply 1.0 to 2.0 gallons of water per acre. For control of plum stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of plum stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

CHERRIES: For control of cherry fruit fly, apply 1.0 to 2.0 gallons of water per acre. For control of cherry stem sawfly, apply 1.0 to 2.0 gallons of water per acre. For control of cherry stem sawfly, apply 1.0 to 2.0 gallons of water per acre.

PEARL ONIONS: For control of pearl onion maggot, apply 1.0 to 2.0 gallons of water per acre. For control of pearl onion maggot, apply 1.0 to 2.0 gallons of water per acre.

SPINACH: For control of spinach leaf miner, apply 1.0 to 2.0 gallons of water per acre. For control of spinach leaf miner, apply 1.0 to 2.0 gallons of water per acre.

CRUCIFEROUS: For control of cabbage root fly, apply 1.0 to 2.0 gallons of water per acre. For control of cabbage root fly, apply 1.0 to 2.0 gallons of water per acre. For control of cabbage root fly, apply 1.0 to 2.0 gallons of water per acre.

Manufactured By

CENTRAL CHEMICAL CORPORATION
General Offices Hagerstown, Maryland

NET CONTENTS 5 GALLONS