

en them to be physically compatible.

# DANGER

## KEEP OUT OF REACH OF CHILDREN

### ANTIDOTES

**IF SWALLOWED:** Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately!

**IF ON SKIN:** Wash immediately with soap and water.

# DANGER



Poisonous by swallowing, inhalation, or skin contact! Do not get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. In case of contact, immediately remove all contaminated clothing and flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention. Wear clean synthetic rubber gloves and a mask or respirator of a type

passed by the U. S. Department of Mines for endrin protection. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed and foodstuffs.

This product is poisonous to fish and wildlife.

Birds feeding on treated areas may be killed.

Keep out of lakes, streams or ponds.

Do not apply to areas where runoff or flooding will contaminate ponds, lakes or streams, or domestic water supplies.

Do not apply when weather conditions favor drift from areas treated.

### RECOMMENDATIONS

**APPLE ORCHARDS:** Pine Mice — Boom Application — use 3 pints of Endrin 1.6 E per 100 gallons of water and apply 100 gallons of spray in a band eleven feet wide and 670 feet long on each side of a tree row (350 to 400 gallons per acre). Hand Gun Application — use 3 pints of Endrin 1.6 E per 100 gallons of water and apply 11 gallons of spray to each under tree area (5½ gallons per side). In order to penetrate ground cover to reach pine mice trail system, apply all sprays at 500 psi pressure. Specific directions concerning nozzles should be obtained from local authorities. Do not disturb trail systems by cultivation before spraying. Allow a minimum of two months before cultivating after applying spray. Make post-harvest applications during October or November in accordance with local recommendations.

Meadow Mice — use 1 quart of Endrin 1.6 E per 100 gallons of water and apply 300 to 350 gallons of spray per acre as directed for pine mice control using hand gun or boom sprayers to wet ground cover under trees. Make post-harvest applications in October and November in accordance with local recommendations.

**POST ALL TREATED AREAS.** Prevent entry to treated area until 30 days after treatment. Do not graze orchards, cut forage for hay or allow drop fruit to be utilized for any purpose.

Do not reuse empty drum. Destroy by perforating or crushing and burying in a safe place.

MANUFACTURED BY

**CENTRAL CHEMICAL CORPORATION**  
General Offices \_\_\_\_\_ GALLONS NET Hagerstown, Maryland

EPA Reg. no. 477-288